arXiv:1304.7527v1 [math.DG] 28 Apr 2013

SINGULAR SUPERSPACES

ALEXANDER ALLDRIDGE, JOACHIM HILGERT, AND TILMANN WURZBACHER

Abstract. We introduce a wide category of superspaces, called locally finitely generated, which properly includes supermanifolds but enjoys much stronger permanence properties, as are prompted by applications. Namely, it is closed under taking finite fibre products (*i.e.* is finitely complete) and thickenings by spectra of Weil superalgebras. Nevertheless, in this category, morphisms with values in a supermanifold are still given in terms of coordinates. This framework gives a natural notion of relative supermanifolds over a locally finitely generated base. Moreover, the existence of inner homs, whose source is the spectrum of a Weil superalgebra, is established; they are generalisations of the Weil functors defined for smooth manifolds.

Introduction

Supermanifolds form a natural $\mathbb{Z}/2\mathbb{Z}$-graded generalisation of the category of manifolds, originally introduced as a tool to treat fermionic and bosonic fields on the same footing. In stark contrast to the traditional theory of smooth manifolds, certain basic constructions in supergeometry automatically lead to 'singular' superspaces. One is thus compelled to work within a wider category.

Indeed, let us make the reasons for this step more explicit. Typically, fibre products of supermanifolds do not exist as such. For instance, assume given the action of a Lie supergroup G on a supermanifold X and a generalised point $x : T \to X$. If $T \neq *$, *e.g.* $T = \mathbb{A}^{0|1}$, that is, the case of an 'odd point', then the isotropy of G at x should be defined as a fibre product, which however usually is not a supermanifold.

Furthermore, as applications in *e.g.* Lie theory or variational calculus suggest, it is natural to consider a category stable under thickening by spectra of finite-dimensional local superalgebras. These are the so-called Weil superalgebras, and include examples such as the dual and super-dual numbers, the jet algebras, and the superpoints. This point of view also gives a natural construction of the 'extensions' of (super) manifolds introduced by A. Weil (and called Weil functors here), as inner hom functors in this category.

Another motivation comes from the necessity for the definition of supermanifolds over a general base, which itself need not be a supermanifold. Notably, this is required for a full-blown theory of Berezin integration along the fibres.

The aim of the present article is to construct an extension of the category of supermanifolds, closed under finite limits and Weil thickenings, which we now proceed to describe.

In algebraic geometry, the objective of constructing a category that meets these requirements is attained by the introduction of the concept of schemes, including *ab initio* singular, *i.e.* non-smooth, objects. Since their inception in the work of

2010 *Mathematics Subject Classification.* Primary 58A50; Secondary 14M30, 32C11.

Key words and phrases. inner hom, fibre product, relative supermanifold, superspace, Weil functor, Weil superalgebra.

We would like to thank SFB/TR 12 "Symmetries and University in Mesoscopic Systems", funded by Deutsche Forschungsgemeinschaft (DFG). The first named author gratefully acknowledges support received from the Leibniz prize awarded to M. Zirnbauer, DFG grant no. ZI 513/2-1.

Berezin, Kostant, and Leites, supermanifolds have been defined within the larger category of superspaces, *i.e.* locally super-ringed spaces, and thus, one may attempt to find a larger subcategory in which the above difficulties are no longer present. In this paper, we single out the full subcategory of *locally finitely generated* superspaces. Here, a superspace is called locally finitely generated if it has a cover by open subspaces, each of which admits an embedding into (smooth or analytic) affine superspace $\mathbb{A}^{m|n}$.

Morally, the local finite generation of a superspace is analogous to the condition that a scheme be *locally of finite type*. However, the ring of germs of $\mathcal{C}^\infty$ functions on $\mathbb{R}^n$ is far from being Noetherian, so the study of these spaces is not merely a translation of methods from algebraic geometry, but requires different techniques.

Indeed, for the embeddings in the definition of locally finite generation, we are obliged to require the property of *tidiness*. Technically, this means that their vanishing ideals satisfy the Whitney condition [9], which in the smooth case is equivalent to closedness of the ideal in the $\mathcal{C}^\infty$ topology (although this is not the form in which it is applied here). Without the condition of tidiness, the existence of equalisers would not be given. Moreover, in their construction, tidiness has to be enforced by a univeral construction, which we call *tidying*.

Another respect, in which we have to depart from the paradigm of algebraic geometry, concerns the description of morphisms to affine superspace $\mathbb{A}^{m|n}$. Morphisms from any locally ringed space to an affine scheme are in one-to-one correspondence with morphisms of the algebras of global sections, a fact which is quickly deduced by localisation. In particular, morphisms to *algebraic* affine space $\mathbb{A}^m$ are given by coordinates; that is, they are in bijection with tuples of functions $(x_1, \dots, x_m)$.

By contrast, to determine to which extent morphisms with range in *smooth* (or analytic) affine superspace $\mathbb{A}^{m|n}$ can be described by coordinates is a subtle matter. For the case of supermanifolds, the fact that they can is a fundamental result due to Leites [7]. We call a superspace for which this theorem holds for all open subspaces *Leites regular* and investigate the stability of this property under passage to colimits, locally closed subspaces, and thickenings. An important outcome of this study is that tidiness is crucial to ensure the hereditarity of Leites regularity; in particular, locally finitely generated superspaces are Leites regular.

An important aspect of Leites regularity is that in the category of Leites regular superspaces, the smooth (or analytic) affine superspace $\mathbb{A}^m$ is the m-fold product of $\mathbb{A}^1$. This is not true in the larger category of superspaces, for the algebra of smooth functions on $\mathbb{R}^2$ is not the tensor square of the ring of univariate smooth functions. Thus, unless one is willing to work with sheaves of topological vector spaces, there is no simple-minded construction of the products $\mathbb{A}^1 \times \cdots \times \mathbb{A}^1$. Instead, we construct smooth (and analytic) affine superspace $\mathbb{A}^m$ directly. The general case $\mathbb{A}^{m|n}$ appears as a special case of the process of *Weil thickening*, by which a superspace is augmented by a Weil superalgebra spectrum.

By tidying, the existence of products for affine superspaces carries over to locally finitely generated superspaces. Together with the existence of equalisers (which exist for locally Hausdorff tidy superspaces), we arrive by the result that the category of locally finitely generated superspaces is finitely complete, contains supermanifolds as a subcategory closed under finite products, and is stable under Weil thickening.

This setup allows for the introduction of *relative supermanifolds* over a locally finitely generated base. Moreover, the *Weil functors* (also known as *Weil extensions* in the literature) and relative versions thereof can be defined as inner hom functors in the category of locally finitely generated superspaces. We show that they are

representable by relative supermanifolds. This gives a uniform construction of the (relative) even and odd tangent bundles, and their higher analogues. Moreover, it has important applications in algebraic topology, notably in the work of Stolz and Teichner on Euclidean field theories, where these inner homs play a prominent role, see *e.g.* Ref. [10].

We end this introduction by giving a brief synopsis of the article's contents; herein, the reader will find references to its quintessential concepts and results, as alluded to above. In Section 1, we recall standard notions from the theory of superspaces. In Section 2, we embark upon a study of the fine structure of embeddings (called immersions in algebraic geometry). In particular, we introduce the notion of *girth* (Definition 2.1), which quantifies the relative size of an ambient space around an embedding. This is crucial below, to prove that Leites regularity is hereditary under certain embeddings. Important examples of embeddings are furnished by the *Weil thickenings*, which are also introduced in this section. Finally, we discuss the concept of tidiness and show how it can be imposed on an embedding by a process called *tidying* (Proposition 2.34).

In Section 3, we define several variants of the standard affine superspace $\mathbb{A}^{m|n}$: real smooth, real analytic, complex analytic, and real smooth and analytic versions with complexified function sheaves. The latter two serve as local models for J. Bernstein's category of *cs* manifolds. We then define *Leites regular* superspaces; this framework gives meaning to the concept of *local coordinate systems*.

In Section 4, we show that Leites regularity is stable under under colimits (Proposition 4.4) and investigate its stability with respect to embeddings (Proposition 4.10) and thickenings (Proposition 4.13), the latter of which Weil thickenings are an example of. We then introduce locally finitely generated superspaces. We show that these are Leites regular (Proposition 4.16), admit finite limits (Corollary 4.23), and Weil thickenings. In this framework, we introduce *relative supermanifolds* over a base, which is locally finitely generated. At this level, base change becomes meaningful, and for a fixed base, the relative category enjoys the same properties as the category of usual supermanifolds. Finally, we introduce (relative) *Weil functors* as inner homs with respect to the spectra of Weil superalgebras, and prove their representability.

Acknowledgements. We extend our thanks to the following institutions for providing their stimulating research environments during the preparation of this article: Mathematisches Forschungsinstitut Oberwolfach, Max-Planck-Institut für Mathematik Bonn, and Ruhr-Universität Bochum. We thank Torsten Wedhorn for useful comments on a preliminary version of the manuscript.

1. Generalities on superspaces

In this section, we collect basic facts and definitions related to superspaces in the sense of Ref. [8]. Most of these are more or less straightforward generalisations of the ungraded case, *v.* Ref. [4]. Accordingly, we will only give full proofs in such cases where we deviate from the standard lore.

1.1. **Superspaces and their morphisms.** We start with the definition of the category of super-ringed spaces.

Definition 1.1 (Super-ringed spaces)**.** A *super-ringed space* is a pair $X = (X_0, \mathcal{O}_X)$, where X_0 is a topological space and $\mathcal{O}_X$ is a sheaf with values in the symmetric monoidal category of supercommutative unital superrings (*i.e.* $\mathbb{Z}/2\mathbb{Z}$ graded rings) and even, unital ring morphisms. It is called the *structure sheaf* of X.

By definition, a *morphism of super-ringed spaces* $\varphi : X \to Y$ is a pair $(\varphi_0, \varphi^\sharp)$ where $\varphi_0 : X_0 \to Y_0$ is a continuous map and $\varphi^\sharp : \varphi_0^{-1}\mathcal{O}_Y \to \mathcal{O}_X$ is a morphism of sheaves (of superrings). These form a category denoted by **SRSp**.

Definition 1.2 (Open subspaces)**.** Whenever X is a super-ringed space and $U \subseteq X_0$ is an open subset, let $X|_U = (U, \mathcal{O}_X|_U)$. Such a super-ringed space is called an *open subspace* of X. It comes with a morphism $j_{X|_U} : X|_U \to X$, defined by $j_{X|_U} \coloneqq (j_{X|_U,0}, j^\sharp_{X|_U})$ where $j_{X|_U,0}$ is the inclusion of U in X_0, and the sheaf morphism $j^\sharp_{X|_U} : j^{-1}_{X|_U,0}\mathcal{O}_X = \mathcal{O}_X|_U \to \mathcal{O}_X|_U$ is the identity; $j_{X|_U}$ is called the *inclusion* of $X|_U$ in X. We will systematically identify open subspaces and the underlying open sets. Thus, unions and finite intersections of open subspaces make sense.

If the sections of the structure sheaf are to represent 'superfunctions' and non-vanishing 'superfunctions' are to be invertible one needs the stalks of the structure sheaf to be local rings. Ringed spaces with this property are called *locally ringed spaces* in algebraic geometry. Recall that a ring R is called *local* if it possesses a unique maximal ideal. Similarly, one says that a superring R is *local*, if it possesses a unique maximal graded ideal. It turns out that in the supercommutative case, one does not have to distinguish between maximal ideals and maximal graded ideals.

Proposition 1.3 (Local superrings)**.** *Let R be a supercommutative superring. The following are equivalent:*

(i) *R is local as a superring;*
(ii) *R is local as an ungraded ring;*
(iii) *for homogeneous $a, b \in R$ with $a + b = 1$, one of a and b is invertible; and*
(iv) *for any $a, b \in R$ such that $a + b = 1$, one of a and b is invertible.*

In this case, the unique maximal ideal is graded and consists of those elements of R which are not invertible.

In view of Proposition 1.3, the following definition of the category of superspaces is an extension of the definition of locally ringed spaces.

Definition 1.4 (Superspaces)**.** A super-ringed space $X = (X_0, \mathcal{O}_X)$ is a *superspace* if for each $x \in X_0$, the stalk $\mathcal{O}_{X,x}$ is a local superring. We denote the maximal ideal by $\mathfrak{m}_{X,x}$. A morphism $\varphi : X \to Y$ of super-ringed spaces where X and Y are superspaces is called *local* if $\varphi^\sharp(\mathfrak{m}_{Y,\varphi_0(x)}) \subseteq \mathfrak{m}_{X,x}$. The category of superspaces and local morphisms is denoted by **SSp**. The locally defined sections of $\mathcal{O}_X$, where X is a superspace, are called *superfunctions*.

Given a field k, a superspace X is called a *k-superspace* if $\mathcal{O}_X$ is a sheaf of k-algebras. The category of k-superspaces and local morphisms φ such that $\varphi^\sharp$ is k-linear is denoted by $\mathbf{SSp}_k$.

There is also a relative version of the category of superspaces.

Definition 1.5 (Relative superspaces)**.** Let $p : X \to S$ be a morphism of superspaces. We will say that X is a *superspace over S* (or a *relative superspace*) and write X/S, the morphism $p = p_X$ then being understood.

A *morphism over S*, written $f : X/S \to Y/S$, is a morphism $f : X \to Y$ such that $p_Y \circ f = p_X$, *i.e.* the following diagram commutes:



We denote the set of morphisms $X \to Y$ over S by $\mathrm{Hom}_S(X, Y)$ and the category of superspaces over a superspace S by $\mathbf{SSp}_S$.

More generally, given any morphism $h : S \to T$, a morphism $f : X \to Y$, where X/S and Y/T are relative superspaces, is *over* h if $p_Y \circ f = h \circ p_X$. This enables us to consider relative superspaces over varying bases.

Remark 1.6 (Terminal objects). The category of k-superspaces has the terminal object $\operatorname{Spec} k \coloneqq (*, k)$. Any k-superspace may therefore be considered as a superspace over $(*, k)$. In fact, the following holds.

Proposition 1.7. *Let k be a (purely even) field. Given a superspace X, there is a bijection between the k-superspace structures on X and morphisms $X \to \operatorname{Spec} k$. Moreover, if X and Y are k-superspaces, and $\varphi : X \to Y$ is a morphism of superspaces, then φ is a morphism of k-superspaces if and only if it is over* $\operatorname{Spec} k$.

We end this subsection with a simple but fundamental construction.

Construction 1.8 (Change of base field)**.** Let ℓ be a field extension of the field k. Any ℓ-superspace X can be naturally considered as a k-superspace by forgetting the ℓ-structure on $\mathcal{O}_X$. In other words, by composing the given morphism $X \to \operatorname{Spec} \ell$ of superspaces with the natural morphism $\operatorname{Spec} \ell \to \operatorname{Spec} k$. We will denote the k-superspace associated with X by the same letter.

Conversely, let X be a k-superspace. We define X_ℓ to be the ℓ-superspace

$$X_\ell \coloneqq (X_0, \mathcal{O}_X \otimes_k \ell).$$

By considering the sheaf embedding $\mathcal{O}_X \to \mathcal{O}_{X_\ell}$ given by construction, we obtain a morphism $X_\ell \to X$ of k-superspaces. Moreover, there is a natural morphism $X_\ell \to \operatorname{Spec} \ell$ of k-superspaces. Let Y be a k-superspace, and assume given morphisms $\varphi : Y \to X$ and $Y \to \operatorname{Spec} \ell$ of k-superspaces. The latter means that Y has an ℓ-superspace structure compatible with the embedding $k \subseteq \ell$. Thus, $\varphi^\sharp$ extends to a unique ℓ-linear map $\mathcal{O}_X \otimes_k \ell \to \varphi_{0,*}\mathcal{O}_Y$. This shows that X_ℓ has the universal property of the fibre product $X \times_{\operatorname{Spec} k} \operatorname{Spec} \ell$ in $\mathbf{SSp}_k$. Thus, the functor $(-)_\ell$ is right adjoint to the forgetful functor $\mathbf{SSp}_\ell \to \mathbf{SSp}_k$.

1.2. **Embeddings.** In this subsection, we discuss a particular class of morphisms, which we call embeddings, following the standard terminology of differential geometry. In algebraic geometry, they are called *immersions*. However, the immersions of differential geometry in general are neither injective, nor is their image locally closed. To avoid confusions in the applications to supermanifolds, we therefore stick to the wording common in the study of smooth manifolds.

Recall that a continuous map is called an *embedding* if it induces a homeomorphism onto its image (endowed with the relative topology). An embedding is called *open* resp. *closed* if its image is, in addition, open resp. closed. An open (resp. closed) embedding is indeed an open (resp. closed) map.

Definition 1.9 (Embeddings)**.** Let $\varphi : Y \to X$ be a morphism of superspaces. It is a called an *open embedding* if it factors as $\varphi = j_{X|_U} \circ \psi$ where $U \subseteq X_0$ is an open subset, and $\psi : Y \to X|_U$ is an isomorphism of superspaces. In this case, $U = \varphi_0(Y_0)$, and we denote by $\varphi(Y)$ the open subspace $X|_U \subseteq X$.

The morphism φ is called a *closed embedding* if the map φ_0 is a closed embedding, and the sheaf morphism $\varphi^\sharp : \mathcal{O}_X \to \varphi_{0,*}\mathcal{O}_Y$ is surjective, *i.e.* it induces an isomorphism

$$\mathcal{O}_X/\mathcal{I}_Y \to \varphi_{0,*}\mathcal{O}_Y,$$

where $\mathcal{I}_Y \coloneqq \ker \varphi^\sharp$ is by definition the *vanishing ideal* of Y (or of φ).

Moreover, φ is called an *embedding* if it factors as $\varphi = \varphi'' \circ \varphi'$ where φ' is a closed embedding and φ'' is an open embedding. Finally, an embedding φ is called a *thickening* if $X_0 = Y_0$ as topological spaces and φ_0 is the identity map.

Open and closed embeddings are embeddings, and embeddings are preserved under field extension. Compositions of open (resp. closed) embeddings are open (resp. closed) embeddings. General embeddings are stable under composition, as follows for instance from the following proposition.

Proposition 1.10. *Let $\varphi : Y \to X$ be a morphism of superspaces. The following assertions are equivalent:*

(i) *The morphism φ is an embedding.*

(ii) *The map φ_0 is an embedding, and there exists an open subset $U \subseteq X_0$ such that $\varphi_0(Y_0)$ is closed in U and $\varphi^\sharp : \mathcal{O}_X|_U \to \varphi_{0,*}\mathcal{O}_Y$ is surjective.*

(iii) *The map φ_0 is an embedding, $\varphi_0(Y_0)$ is locally closed, and the sheaf map $\varphi^\sharp : \mathcal{O}_X|_U \to \varphi_{0,*}\mathcal{O}_Y$ is surjective, where $U \coloneqq \varphi_0(Y_0) \cup X_0 \setminus \overline{\varphi_0(Y_0)}$.*

The proposition is implied by the following basic topological fact.

Lemma 1.11. *Let T be a topological space and S a locally closed subset. The largest open subset $U \subseteq T$ such that S is closed in U is $U \coloneqq S \cup T \setminus \overline{S}$.*

The characterisation of embeddings in Proposition 1.10 also allows us to decide when a morphism factors through an embedding.

Proposition 1.12 (Factorisation through embeddings)**.** *Let $\varphi : Y \to X$ be an embedding with ideal $\mathcal{I}_Y \subseteq \mathcal{O}_X|_U$, where $U \coloneqq \varphi_0(Y_0) \cup X_0 \setminus \overline{\varphi_0(Y_0)}$, and $\psi : Z \to X$ a morphism. Then ψ factors through φ if and only if $\psi_0(Z_0) \subseteq \varphi_0(Y_0)$ and $\mathcal{I}_Y \subseteq \ker \psi^\sharp|_U$. This factorisation is unique.*

If the map underlying an embedding is an inclusion we talk about a subspace.

Definition 1.13 (Subspaces and closed subspaces)**.** Let X be a superspace. If $j : Y \to X$ is a (closed) embedding of superspaces where $Y_0 \subseteq X_0$ and j_0 is the inclusion of Y_0 in X_0, then we say that Y is a (closed) *subspace* of X.

Given a locally closed subset Y_0 of X_0, the structure of a subspace Y over Y_0 (if it exists) is determined uniquely up to canonical isomorphism by a graded ideal $\mathcal{I}_Y$ of $\mathcal{O}_X|U$, where $U \coloneqq Y_0 \cup X_0 \setminus \overline{Y_0}$, such that $\operatorname{supp} \mathcal{O}_X|_U/\mathcal{I}_Y = Y_0$. This ideal is called the *vanishing ideal* of Y. The isomorphism classes of such subspaces Y are parametrised by the ideals $\mathcal{I}_Y$.

For any subspace Y of a superspace X there is an increasing sequence of subspaces all with underlying space Y_0. The sections of the corresponding structure sheaves should be viewed as superfunctions on X which do not necessarily *vanish* on Y_0, but are *nilpotent* of a certain order on Y_0.

Definition 1.14 (Infinitesimal normal neighbourhoods)**.** Let $j : Y \to X$ be an embedding of superspaces with ideal $\mathcal{I}_Y \subseteq \mathcal{O}_X|_U$, where $U \coloneqq j_0(Y_0) \cup X_0 \setminus \overline{j_0(Y_0)}$. For any integer $n \geqslant 0$, let $\mathcal{I}_{Y^{(n)}} \coloneqq \mathcal{I}_Y^{n+1} \subseteq \mathcal{O}_X|_U$ and $\mathcal{O}_{Y^{(n)}} \coloneqq j_0^{-1}(\mathcal{O}_X|_U/\mathcal{I}_{Y^{(n)}})$.

The superspace $Y^{(n)} \coloneqq (Y_0, \mathcal{O}_{Y^{(n)}})$ is called the *nth infinitesimal normal neighbourhood* of Y in X. It comes with a natural embedding $j^{(n)} : Y^{(n)} \to X$. In particular, $Y = Y^{(0)}$ and $j = j^{(0)}$. Whenever $n \leqslant m$, the embedding $j^{(n)} : Y^{(n)} \to X$ factors uniquely through $j^{(m)} : Y^{(m)} \to X$ *via* a closed embedding $j^{(nm)} : Y^{(n)} \to Y^{(m)}$, by virtue of Proposition 1.12.

The $(Y^{(n)}, j^{(nm)})$ form an inductive system of superspaces, so naturally the question arises whether we can form inductive limits in $\mathbf{SRSp}_S$ and $\mathbf{SSp}_S$. It turns out that not only inductive limits but *all* small colimits exist in these categories.

Proposition 1.15. *Let S be a super-ringed space. The category $\mathbf{SRSp}_S$ is cocomplete,* i.e. *all small colimits exist, and $\mathbf{SSp}_S$ is closed under all of them.*

Proof. By Proposition 1.3, superrings are local if and only if they are local as ungraded rings, so the statement follows from the classical case [3, I.1.6]. □

The Yoneda embedding allows us to identify the structure sheaves of small colimits of superspaces.

Remark 1.16. Let $X = \varinjlim_k X_k$ be an inductive limit in $\mathbf{SRSp}_S$ (or in $\mathbf{SSp}_S$). Then $\mathcal{O}_X = \varprojlim_k (f_k)_{0,*}\mathcal{O}_{X_k}$ in $\mathbf{Sh}(X_0)$, where $f_k : X_k \to X$ are the natural morphisms. Because this projective limit is the projective limit of presheaves and the Yoneda embedding commutes with projective limits, we have that $\Gamma(\mathcal{O}_X) = \varprojlim_k \Gamma(\mathcal{O}_{X_k})$.

The statement also holds with 'small colimit' in place of 'inductive limit' if one replaces 'projective limit' by 'small limit'.

Example 1.17. We return to the inductive system of infinitesimal normal neighbourhoods. Let $Y^{(\infty)}$ with the morphisms $j^{(n\infty)} : Y^{(n)} \to Y^{(\infty)}$ denote the inductive limit of the inductive system $(Y^{(n)}, j^{(nm)})$ of superspaces. The superspace $Y^{(\infty)}$ is called the *infinitesimal normal neighbourhood of order* ∞ of Y in X. The morphisms $j^{(nm)}$, where $m \in \mathbb{N} \cup \infty$, are thickenings.

1.3. **Gluing.** The gluing construction is standard in geometry. It depends on the possibility to have fibre products with open embeddings.

Proposition 1.18 (Fibre products with open embeddings)**.** *Let $\varphi : X \to Z$ be an open embedding of superspaces, and $\psi : Y \to Z$ a morphism of superspaces. Then $X \times_Z Y$ exists in* $\mathbf{SSp}$*, and $p_2 : X \times_Z Y \to Y$ is an open embedding. The statement carries over to k-superspaces.*

Now we can show how to glue superspaces and their morphisms; this technique will be fundamental. To have it available in its most general form, we introduce the concept of 'gluing data'.

Definition 1.19 (Coverings and gluing data)**.** Let S be a superspace. An *open cover of S* is a family (U_i) of open subspaces such that $\bigcup_i U_{i,0} = S_0$.

More generally, a morphism $\varphi : T \to S$ is called a *covering of S* if there exists an open cover (U_i) of T such that $T = \coprod_i U_i$, $\varphi|_{U_i} : U_i \to S$ are open embeddings, and the open subspaces $\varphi(U_i) \subseteq S$ form an open cover of S. In other words, $\varphi = \coprod_i \varphi_i$ where $\varphi_i : U_i \to S$ are open embeddings. We identify φ with the collection (φ_i).

Given two coverings $\varphi = (\varphi_i : U_i \to S)$ and $\psi = (\psi_j : V_j \to S)$ of S, the fibre product $\varphi \times_S \psi = (\varphi_i \times_S \psi_j)$ exists, in view of Proposition 1.18. In case $\psi = \varphi$, we write $\varphi_{ij} : U_{ij} \coloneqq U_i \times_S U_j \to S$ for this fibre product. The triple intersections $U_i \times_S U_j \times_S U_k \to S$ will be denoted by $\varphi_{ijk} : U_{ijk} \to S$.

Let $\varphi = (\varphi_i : U_i \to S)$ be a covering of S, and assume given superspaces X_i/U_i and isomorphisms $\psi_{ij} : X_j \times_{U_j} U_{ij} \to X_i \times_{U_i} U_{ij}$. The collection of data $(\varphi_i : U_i \to S, X_i/U_i, \psi_{ij})$ will be called *gluing data for a superspace over S* if

$$p_{12}^*(\psi_{ij}) \circ p_{23}^*(\psi_{jk}) \circ p_{13}^*(\psi_{ki}) = \mathrm{id}_{X_i \times_{U_i} U_{ijk}}. \tag{1.1}$$

Here, $p_{12}^*\psi_{ij}$ is the morphism $X_j \times_{U_j} U_{ijk} \to X_i \times_{U_i} U_{ijk}$ induced by ψ_{ij} by pulling back along the left-hand vertical face of the pullback square

$$\begin{array}{ccc} U_{ijk} & \xrightarrow{p_3} & U_k \\ {\scriptstyle p_{12}}\downarrow & & \downarrow{\scriptstyle \varphi_k} \\ U_{ij} & \xrightarrow[\varphi_{ij}]{} & S \end{array}$$

Similarly for the other quantites in Equation (1.1).

When the U_i are simply open subspaces of S and the φ_i are just the canonical embeddings $j_{U_i} : U_i \to S$, we will forgo mentioning φ explicitly and simply write

$(U_i, X_i/U_i, \psi_{ij})$ for the gluing data. In this case, somewhat abusing notation, we will write Equation (1.1) in the simpler form

$$\psi_{ij} \circ \psi_{jk} \circ \psi_{ki} = \mathrm{id} \quad \text{on} \quad X_i \times_{U_i} U_{ijk}.$$

Once one has a set of gluing data it follows from the corresponding statements for sheaves that they lead to glued superspaces.

Proposition 1.20. *Let S be a superspace and $(\varphi_i : U_i \to S, X_i/U_i, \psi_{ij})$ gluing data for a superspace over S. Then there exist a superspace X/S and isomorphisms $\psi_i : X_i/U_i \to (X \times_S U_i)/U_i$ with $\psi_j = \psi_i \circ \psi_{ij}$ on $X_j \times_{U_j} U_{ij}$.*

Moreover, these data are uniquely characterised up to unique isomorphism by the following universal property: For any superspace Y/S and any choice of morphisms $\varrho_i : X_i/U_i \to (Y \times_S U_i)/U_i$ such that $\varrho_j = \varrho_i \circ \psi_{ij}$ on $X_j \times_{U_j} U_{ij}$, there exists a unique morphism $\varrho : X/S \to Y/S$ such that $\varrho \circ \psi_i = \varrho_i$.

The statement of Proposition 1.20 immediately carries over to k-superspaces.

2. The local structure of embeddings

In this section, we introduce some concepts, which help us in the quantitative study of embeddings. We also define for any superspace certain distinguished subspaces (namely, the reduction and the body), as well as a useful class of thickenings (namely, the Weil thickenings). We analyse the fine structure of embeddings by the use of the new concept of tidiness. Finally, we show how to improve embeddings by a regularisation process called tidying.

2.1. Girth and retractions of embeddings.

There are thickenings of various sizes, as is illustrated by the example of infinitesimal normal neighbourhoods. We introduce the concept of girth of an embedding to quantify this phenomenon. Conversely, one can consider the question whether a thickening can be reversed. This leads to the notion of a retraction.

Definition 2.1 (Finite and countable girth)**.** Let $j_Y : Y \to X$ be an embedding of S-superspaces. We say that X (or j_Y) has *finite girth around Y* if the vanishing ideal $\mathcal{I}_Y$ is nilpotent, *i.e.* if there exists some non-negative integer q such that $\mathcal{I}_Y^{q+1} = 0$. In this case, the minimal q will be called the *girth* of X around Y. We shall say that X (or j_Y) is *locally of finite girth around Y* if there exists an open cover of U (the largest open subset of X_0 in which $j_{Y,0}(Y_0)$ is closed) such that the restriction of $\mathcal{I}_Y$ to every patch of the cover is nilpotent.

Similarly, we say that X (or j_Y) has *countable girth around Y* if $\bigcap_{q=1}^{\infty} \mathcal{I}_Y^q = 0$. If X has locally finite girth around Y, then it has countable girth around Y.

Given an embedding $j : Y \to X$, we obtain for the infinitesimal normal neighbourhoods $Y^{(k)}$: If $m \geqslant n \geqslant 0$ is finite, then the thickening $j^{(nm)} : Y^{(n)} \to Y^{(m)}$ has finite girth at most $\lceil m/n \rceil$.

We have the following useful fact, of which we omit the simple proof.

Proposition 2.2 (Transitivity of girth)**.** *Let $j_Z : Z \to Y$ and $j_Y : Y \to X$ be embeddings of S-superspaces. If X has (locally) finite girth around Y, and Y has (locally) finite girth around Z, then X has (locally) finite girth around Z with respect to the embedding $j_Y \circ j_Z : Z \to X$.*

We now introduce retractions, *i.e.* left inverses for embeddings.

Definition 2.3 (Retractions)**.** Let X be an S-superspace and $j_Y : Y \to X$ a morphism over S which is an embedding. Let $U \subseteq X_0$ be the largest open subset in which $j_{Y,0}(Y_0)$ is closed. A *retraction over Y* is a left inverse $r : X|_U \to Y$ of j_Y, *i.e.* a morphism over S such that the following diagram commutes:



If j_Y admits a retraction, it will be called *retractable*. More generally, if j_Y admits an open cover by retractable embeddings j_{Y_i}, *i.e.* there are covers by open subspaces (Y_i) of Y and (X_i) of X such that the embeddings $j_{Y_i} : Y_i \to X_i$ induced by j_Y are retractable, it will be called *locally retractable*. We shall then also say that X is (locally) retractable around Y.

Let $j_Y : Y \to X$ be an embedding and U the largest open subset of X_0 such that $j_{Y,0}(Y_0)$ is closed in U. Then j_Y is retractable if and only if the short exact sequence

$$0 \longrightarrow \mathcal{I}_Y \longrightarrow \mathcal{O}_X|_U \xrightarrow{j_Y^\sharp} j_{Y_0,*}\mathcal{O}_Y \longrightarrow 0$$

splits as a sequence of superring sheaves. (The splitting is required to be unital on the support $j_{Y,0}(Y_0)$ of $j_{Y_0,*}\mathcal{O}_Y$.) If j_Y is a morphism over S, then it is retractable as such if and only if the sequence splits as a sequence of $p_{X,0}^{-1}\mathcal{O}_S$-algebras.

As in the case of girth, we have the following useful fact.

Proposition 2.4 (Transitivity of retractability)**.** *Let $j_Z : Z \to Y$ and $j_Y : Y \to X$ be embeddings of S-superspaces. If X is (locally) retractable around Y, and Y is (locally) retractable around Z, then X is (locally) retractable around Z with respect to the embedding $j_Y \circ j_Z : Z \to X$.*

Let $j_Y : Y \to X$ be an embedding. If X has girth 0 around Y, then Y is an open subspace of X, so X is retractable around Y.

2.2. **Reduction and body.** Reduction of locally ringed spaces removes nilpotency. Both for schemes and for supermanifolds, this results in a space with a sheaf of functions whose sections are determined completely by their values.

For superspaces, one has yet another reduction, by which everything generated by odd superfunctions is removed. The outcome is called the body of the superspace.

Construction 2.5 (Reduction of superspaces)**.** Let X be a superspace and $x \in X_0$ a point. We define $\varkappa(x) \coloneqq \varkappa_X(x) \coloneqq \mathcal{O}_{X,x}/\mathfrak{m}_{X,x}$; this is a field called the *residue field* at x. If $U \subseteq X_0$ is open, $x \in U$, and $f \in \mathcal{O}_X(U)$, we define $f(x) \in \varkappa(x)$ to be the image of the germ f_x in $\varkappa(x)$. This quantity is called the *value* of f at x.

For any open subset $U \subseteq X_0$, define an ideal $\mathcal{N}_X \subseteq \mathcal{O}_X$ by

$$\begin{aligned}\mathcal{N}_X(U) &\coloneqq \big\{ f \in \mathcal{O}_X(U) \;\big|\; f(x) = 0 \text{ for all } x \in U \big\} \\ &= \big\{ f \in \mathcal{O}_X(U) \;\big|\; f_x \in \mathfrak{m}_{X,x} \text{ for all } x \in U \big\}.\end{aligned}$$

Define $\mathcal{O}_{X_0} \coloneqq \mathcal{O}_X/\mathcal{N}_X$. Endowed with this sheaf, X_0 is a superspace, which by abuse of notation, we will denote by X_0 and call the *reduced superspace* or *reduction* of X. If X is a k-superspace, then so is X_0. In the latter case, $\varkappa(x)$ is an extension of k for any $x \in X_0$. There is a canonical morphism of (k-)superspaces $j_{X_0} : X_0 \to X$, defined by

$$j_{X_0} \coloneqq \big(j_{X,0} = \mathrm{id}_{X_0}, \mathcal{O}_X \to \mathcal{O}_X/\mathcal{N}_X = \mathcal{O}_{X_0}\big)$$

which is by its mere definition a thickening with vanishing ideal $\mathcal{N}_X$. If j_{X_0} is an isomorphism, then X is called *reduced*.

The reduction satisfies a universal property and thus defines a functor.

Lemma 2.6. *Let X be a superspace. Then any morphism $Y \to X$, where Y is reduced, factors uniquely through j_{X_0}.*

Proposition 2.7 (Functoriality of reduction)**.** *There is a* reduction functor*, denoted by $(-)_0 : \mathbf{SSp} \to \mathbf{SSp}$, which is defined on objects by $X \mapsto X_0$, and on morphisms by letting $\varphi_0 : X_0 \to Y_0$ for $\varphi : X \to Y$ be the unique factorisation of $\varphi \circ j_{X_0}$ through j_{Y_0}. It is right adjoint to the inclusion of reduced superspaces in superspaces.*

The canonical morphism $j_{X_0} : X_0 \to X$ defines a natural transformation of functors $(-)_0 \to \mathrm{id}$. The statement carries over to k-superspaces.

We will call a retraction of the embedding $j_{X_0} : X_0 \to X$ simply a *retraction* of X. A retraction is the same as the datum of a sheaf morphism $\mathcal{O}_{X_0} \to \mathcal{O}_X$ of local superrings such that for the induced $\mathcal{O}_{X_0}$-algebra structure on $\mathcal{O}_X$, the canonical morphism $\mathcal{O}_X \to \mathcal{O}_{X_0}$ is $\mathcal{O}_{X_0}$-linear.

In more conventional terminology, if r is a retraction, one says that the image under $r^\sharp$ of $\Gamma(\mathcal{O}_{X_0})$ in $\Gamma(\mathcal{O}_X)$ is a *function factor*. Reductions play an important role in the theory of integration on supermanifolds.

Similarly, we shall say that X has *finite girth*, *girth h*, *locally finite girth*, or *countable girth* if it does so around X_0.

We denote by $\mathcal{N}_X^\infty$ the ideal $\bigcap_{N=1}^\infty \mathcal{N}_X^N$, and call it the *Whitney ideal* of the superspace X. Thus, X has countable girth if and only if $\mathcal{N}_X^\infty = 0$.

Definition 2.8 (Body of a superspace)**.** Let X be a superspace. Define its *body* $X_{\bar 0}$ as the superspace $X_{\bar 0} := (X_0, \mathcal{O}_{X_{\bar 0}})$, where $\mathcal{O}_{X_{\bar 0}} := \mathcal{O}_X/\mathcal{I}_{X_{\bar 0}}$ and $\mathcal{I}_{X_{\bar 0}} := \mathcal{O}_{X,\bar 1} + \mathcal{O}_{X,\bar 1}^2$ is the ideal generated by $\mathcal{O}_{X,\bar 1}$. When X is a k-superspace, then so is $X_{\bar 0}$.

The body is equipped with a thickening $j_{X_{\bar 0}} : X_{\bar 0} \to X$, which is a morphism of k-superspaces if X is a k-superspace. When it is an isomorphism, X is called *even*.

The body of a superspace enjoys analogous properties to the reduction.

Lemma 2.9. *Let X be a superspace. The body $X_{\bar 0}$ enjoys the following universal property: Any morphism $Y \to X$, where Y is even, factors uniquely through $j_{X_{\bar 0}}$.*

Proposition 2.10 (Functoriality of the body)**.** *There is a* body functor*, denoted by $(-)_{\bar 0} : \mathbf{SSp} \to \mathbf{SSp}$, which is defined on objects by $X \mapsto X_{\bar 0}$ and on morphisms by letting $\varphi_{\bar 0} : X_{\bar 0} \to Y_{\bar 0}$ for $\varphi : X \to Y$ be the unique factorisation of $\varphi \circ j_{X_{\bar 0}}$ through $j_{Y_{\bar 0}}$. It is right adjoint to the inclusion of even superspaces in superspaces.*

The canonical morphism $j_{X_{\bar 0}} : X_{\bar 0} \to X$ defines a natural transformation of functors $(-)_{\bar 0} \to \mathrm{id}$. The statement carries over to k-superspaces.

The residue fields in the construction of the reduction allow us to speak about values of 'superfunctions', *i.e.* sections of the structure sheaf in single points of X_0. If X is a k-superspace k is contained in $\varkappa(x)$ for all $x \in X_0$. Thus one can define sections with values in A for any subset A of k.

Definition 2.11 (Superfunctions with specified values)**.** Let $(X, \mathcal{O}_X)$ be a k-superspace. For any $A \subseteq k$, we define a sheaf $\mathcal{O}_{X,A}$ by

$$\mathcal{O}_{X,A}(U) := \big\{ f \in \mathcal{O}_X(U) \;\big|\; f(x) = j_{X_0}^\sharp(f)(x) \in A \text{ for all } x \in U \big\}.$$

The local sections are called *superfunctions with values in A*.

Note that if A is a subfield of k, then $\mathcal{O}_{X,A}$ is an A-superalgebra sheaf. We have the following corollary of Proposition 2.7.

Corollary 2.12. *Let $A \subseteq k$ and $\varphi : X \to Y$ a morphism of k-superspaces. Then $\varphi^\sharp$ induces a sheaf morphism $\varphi_0^{-1}\mathcal{O}_{Y,A} \to \mathcal{O}_{X,A}$.*

Let k be a field and X be a k-superspace. Define

$$X_0(k) \coloneqq \big\{x \in X_0 \;\big|\; \varkappa(x) = k\big\}.$$

Recall that the elements of $X_0(k)$ are called *k-rational points* of X_0. If $X_0 = X_0(k)$, then the local sections of $\mathcal{O}_{X_0}$ may be considered as k-valued functions. In this case, we call X *k-rational*. The following lemma is standard in algebraic geometry.

Lemma 2.13 (k-rational points)**.** *Let $* = (*, k)$ be the terminal object in $\mathbf{SSp}_k$. For any k-superspace X, there is a natural bijection*

$$X(*) \coloneqq \mathrm{Hom}(*, X) \to X_0(k).$$

If $x \in X_0$, then $x \in X_0(k)$ if and only if $\mathcal{O}_{X,x} = k \oplus \mathfrak{m}_{X,x}$.

A morphism between super-ringed spaces which happen to be superspaces is not necessary local, *i.e.* a morphism of superspaces. If the superspaces are k-rational and the ring morphisms k-linear, this is no longer an option.

Proposition 2.14 (Locality of morphisms)**.** *Let X and Y be k-rational k-superspaces and $\varphi : X \to Y$ a morphism of super-ringed space such that $\varphi^\sharp$ is k-linear. Then $\varphi^\sharp(f)(x) = f(\varphi_0(x))$, for any $x \in X_0$ and $f \in \mathcal{O}_{Y,\varphi_0(x)}$. In particular, φ is local, and therefore, a morphism of k-superspaces.*

Proof. Let $y = \varphi_0(x)$. In view of Lemma 2.13 and Proposition 1.3, for $f_y \in \mathcal{O}_{Y,y}$, $f(y) \in \varkappa_Y(y) = k$ is the unique $\lambda \in k$ such that $f_y - \lambda$ is not invertible. By assumption, $\varphi^\sharp_y(f_y) - \lambda = \varphi^\sharp_y(f_y - \lambda)$, and this is invertible whenever $\lambda \neq f(y)$. Let $g = \varphi^\sharp(f)$. Since x is k-rational, there is some $\lambda \in k = \varkappa_X(x)$ for which $g_x - \lambda$ is not invertible, namely, $\lambda = g(x)$. Hence, $\varphi^\sharp(f)(x) = g(x) = f(y)$. In particular, if $f_y \in \mathfrak{m}_{Y,y}$, *i.e.* $f(y) = 0$, then $\varphi^\sharp(f)(x) = 0$, *i.e.* $\varphi^\sharp(f)_x \in \mathfrak{m}_{X,x}$. □

Remark 2.15. In the literature, morphisms of supermanifolds (to be defined below) are often not assumed to be local. Since supermanifolds are rational superspaces, the preceding proposition shows that the locality of morphisms of supermanifolds is automatic. However, in considering morphisms from more general superspaces to supermanifolds, it is more natural to assume locality, as we do.

2.3. **Weil thickenings.** In this subsection, we introduce a construction called Weil thickening, which in its original form goes back to A. Weil [11], compare also Ref. [5]. It does not change the underlying topological space, but only the structure sheaf. Basic examples of superspaces arise by Weil thickening; moreover, it is useful for the infinitesimal study of morphisms. It has two ingredients, a superspace and a Weil superalgebra, which is a special type of local superalgebra.

Definition 2.16 (Weil superalgebras)**.** Let k be a field. A *Weil k-superalgebra* is a finite-dimensional k-superalgebra A with a graded nilpotent ideal $\mathfrak{m}$ such that $A = k \oplus \mathfrak{m}$. Any Weil superalgebra is a local superring, and $\mathfrak{m}$ is the maximal (graded) ideal; it consists exactly of the nilpotent elements of A. The minimal h such that $\mathfrak{m}^{h+1} = 0$ is called the *girth* of A. Equivalently, one may define Weil superalgebras as finite-dimensional local superalgebras (by Krull's theorem).

Weil superalgebras are obtained by truncation of polynomial superalgebras.

Example 2.17. Let A be a k-superalgebra where $\mathrm{char}\, k = 0$. Then A is a Weil superalgebra if and only if $A \cong k[x_1, \dots, x_k|\xi_1, \dots, \xi_\ell]/I$ for some graded ideal I such that $I \supseteq (x_1, \dots, x_k, \xi_1, \dots, \xi_\ell)^N$, for some $N > 0$. In particular, $\bigwedge(k^q)^*$ and $k[\varepsilon]/(\varepsilon^{k+1})$ are Weil k-superalgebras, of girth q and k, respectively.

The Weil thickening X^A of a superspace X by a Weil algebra A is obtained simply by tensoring each ring of sections by A. It turns out that there is a canonical thickening $X \to X^A$ which is a closed embedding and admits a canonical retraction.

Construction 2.18 (Weil thickened superspaces)**.** Let X be a k-superspace. Given a Weil k-superalgebra $A = k \oplus \mathfrak{m}_A$, form $X^A := (X_0, \mathcal{O}_X \otimes A)$, so

$$\mathfrak{m}_{X^A,x} = \mathfrak{m}_{X,x} \otimes A + \mathcal{O}_{X,x} \otimes \mathfrak{m}_A. \tag{2.1}$$

There is a canonical morphism $j_X^A : X \to X^A$, given by $j_{X,0}^A := \mathrm{id}_{X_0}$, and

$$j_X^{A\sharp} := \mathrm{id}_{\mathcal{O}_X} \otimes \varepsilon : \mathcal{O}_{X^A} = \mathcal{O}_X \otimes A \to \mathcal{O}_X \otimes k = \mathcal{O}_X,$$

where $\varepsilon : A \to k$ is the unique algebra morphism. It is clear that j_X^A is a closed embedding, indeed, a thickening. There is a canonical retraction $r_{X^A} : X^A \to X$, given by $r_{X,0}^A := \mathrm{id}_{X_0}$ and

$$r_X^{A\sharp} := \mathrm{id}_{\mathcal{O}_X} \otimes \eta : \mathcal{O}_X = \mathcal{O}_X \otimes k \to \mathcal{O}_{X^A} = \mathcal{O}_X \otimes A,$$

where $\eta : k \to A$ is the unique algebra morphism.

By Proposition 2.4, $r_X^A : X^A \to X$ can be combined with retractions of embeddings of other superspaces into X.

Proposition 2.19 (Retractability of Weil thickenings)**.** *Let $j_Y : Y \to X$ be a retractable embedding of k-superspaces and A a Weil k-superalgebra. Then X^A is retractable around Y.*

Weil thickenings can be reinterpreted as products with superspaces whose underlying topological space is a singleton.

Proposition 2.20. *Let A be a Weil k-superalgebra, and let $\operatorname{Spec} A = (*, A)$. For any k-superspace X, $X \times \operatorname{Spec} A$ exists in $\mathbf{SSp}_k$, and is given by X^A. If B is another Weil k-superalgebra, then $\operatorname{Spec}(A \otimes B) = \operatorname{Spec} A \times \operatorname{Spec} B$.*

Proof. Let Y be a k-superspace. Then $\operatorname{Hom}(Y, \operatorname{Spec} A)$ is the same as the set of all morphisms of k-superalgebras $\psi^\sharp : A \to \Gamma(\mathcal{O}_Y)$. Consider the map

$$\operatorname{Hom}(Y, X) \times \operatorname{Hom}(Y, \operatorname{Spec} A) \to \operatorname{Hom}(Y, X^A) : (\varphi, \psi^\sharp) \mapsto (\varphi_0, \varphi^\sharp \otimes \psi^\sharp),$$

given by $(\varphi^\sharp \otimes \psi^\sharp)(f \otimes a) = \varphi^\sharp(f) \cdot \psi^\sharp(a)$. It is clearly a natural bijection. This proves that $X^A = X \times \operatorname{Spec} A$, and the second statement follows. □

Proposition 2.20 shows that $j_X^A : X \to X^A = X \times \operatorname{Spec} A$ is just $j_X^A = (\mathrm{id}_X, *)$, where $* \to \operatorname{Spec} A$ is the canonical embedding. By the same token, the morphism $r_X^A : X^A = X \times \operatorname{Spec} A \to X$ is just $r_X^A = p_1$, the first projection.

Remark 2.21. In algebraic geometry, one defines $\operatorname{Spec} R$ for any commutative ring R to be the collection of prime ideals with the Zariski topology, endowed with the structure sheaf obtained by localisation of R.

Note that for a Weil k-superalgebra A any prime ideal is maximal, justifying our above definition. Indeed, let $\mathfrak{p}$ be a prime ideal in A. We have $A = k[x_a]/I$ for some set (x_a) of homogeneous indeterminates, and some ideal I containing $(x_a)^N$ for some N. Then $\mathfrak{p} = \mathfrak{q}/I$ for some prime ideal $\mathfrak{q}$ of $k[x_a]$ containing I. This implies $x_a^N \in \mathfrak{q}$ for all a, but since $\mathfrak{q}$ is a radical ideal, it follows that $\mathfrak{q} = (x_a)$, so that $\mathfrak{p} = \mathfrak{m}_A$.

Using Proposition 2.2, we obtain that for any finite girth embedding $j_Y : Y \to X$ the embedding $j_X^A \circ j_Y : Y \to X^A$ into a Weil thickening X^A of X has finite girth. We can, however, do better and determine the girth of j_X^A precisely.

Proposition 2.22 (Girth of Weil thickenings)**.** *Let $j_Y : Y \to X$ be an embedding of k-superspaces of girth q and A a Weil k-superalgebra of girth h.*

Then the embedding $j := j_X^A \circ j_Y : Y \to X^A$ has girth $q + h$. In particular, $X \times \operatorname{Spec} A = X^A$ has girth h around $j_X^A : X \to X^A$. Similarly, if j_Y has countable girth, then so has j.

Proof. We may assume that $j_Y : Y \to X$ is a closed embedding. The vanishing ideal of $j_X^A : X \to X^A$ is simply $\mathcal{O}_X \otimes \mathfrak{m}$, where $\mathfrak{m} = \mathfrak{m}_A$ is the maximal ideal of A. Moreover, by definition of j_X^A, we have

$$j^\sharp(f \otimes a) = j_Y^\sharp(f) \cdot \varepsilon(a) \quad \text{for all} \quad f \otimes a \in \mathcal{O}_{X^A}(U) = \mathcal{O}_X(U) \otimes A.$$

Since $\varepsilon(a) \in k$ and k has no zero divisors, we have $j^\sharp(f \otimes a) = 0$ if and only if $j_Y^\sharp(f) = 0$ or $\varepsilon(a) = 0$. It follows that the vanishing ideal $\mathcal{I}$ of j is given by

$$\mathcal{I} = \mathcal{I}_Y \otimes A + \mathcal{O}_X \otimes \mathfrak{m}$$

where $\mathcal{I}_Y$ is the vanishing ideal of $j_Y : Y \to X$. Next, we compute inductively that

$$\mathcal{I}^j/\mathcal{I}^{j+1} = \textstyle\bigoplus_{i=0}^j \mathcal{I}_Y^i/\mathcal{I}_Y^{i+1} \otimes \mathfrak{m}^{j-i}/\mathfrak{m}^{j-i+1},$$

where we agree to write $\mathcal{I}^0 = \mathcal{O}_{X^A}$, $\mathcal{I}_Y^0 = \mathcal{O}_X$, and $\mathfrak{m}^0 = A$. Therefore, the left hand side is zero if and only $j > q + h$.

If j has countable girth, we argue similarly. Indeed, for $N > h$, we have

$$\mathcal{I}^N \subseteq \textstyle\sum_{k=0}^h \mathcal{I}_Y^{N-k} \otimes \mathfrak{m}^k,$$

so that $\mathcal{I}^N \subseteq \mathcal{I}_Y^{N-h} \otimes A$. Let $f \in \bigcap_{N=1}^\infty \mathcal{I}^N(U)$. Choosing a k-basis (a_ℓ) of A, we have a unique representation $f = \sum_\ell f_\ell \otimes a_\ell$ where $f_\ell \in \mathcal{O}_X(U)$. By the above, $f_\ell \in \bigcap_{N=1}^\infty \mathcal{I}_Y^N(U) = 0$, which proves our claim. □

2.4. **Noetherian jet spaces and tidiness.** In this subsection we introduce two regularity conditions, one for superspaces and one for embeddings of superspaces. Given a superspace X we define a ring of 'jets' for any $x \in X_0$. If these rings are all Noetherian, the sections of the structure sheaf are not too far from being determined by their restrictions to infinitesimal normal neighborhoods of points.

An embedding $j : Y \to X$ of superspaces is called *tidy* if the restriction of $j^\sharp$ to the stalk at some point $x \in j_0(Y_0)$ is determined by the jets in a neighborhood of x in $j_0(Y_0)$. A superspace with Noetherian rings of jets for which the identity is tidy will be call a tidy superspace. The concept of tidiness will be instrumental in the description of superspaces in terms of coordinates.

In this subsection we describe a number of general properties of tidy embeddings and superspaces, starting with some basic facts from commutative algebra which will help us to deal with finiteness properties of the jet spaces.

Let R be a local supercommutative superring with maximal ideal $\mathfrak{m}$. We denote by $\mathfrak{m}^\infty := \bigcap_{N=1}^\infty \mathfrak{m}^N$ the Whitney ideal of the superspace $(*, R)$.

Proposition 2.23 (Krull's theorem)**.** *Assume that $R/\mathfrak{m}^\infty$ is Noetherian. For any graded ideal $I \subseteq R$, we have $\mathfrak{m}^\infty + I = \bigcap_{N=1}^\infty (\mathfrak{m}^N + I)$.*

Proof. Since $R/\mathfrak{m}^\infty$ is local and Noetherian, so is the superring $S := R/(I + \mathfrak{m}^\infty)$. The maximal ideal of S is $\mathfrak{n} := \mathfrak{m}/(\mathfrak{m}^\infty + I)$. Since S is Noetherian, $\mathfrak{n}^\infty$ is finitely generated, and $\mathfrak{n} \cdot \mathfrak{n}^\infty = \mathfrak{n}^\infty$. Hence, by the Nakayama Lemma, $\mathfrak{n}^\infty = 0$. But then $\bigcap_{N=1}^\infty (\mathfrak{m}^N + I) = \mathfrak{m}^\infty + I$, as desired. □

Definition 2.24 (Noetherian jet spaces)**.** Let X be a superspace. If $\mathcal{O}_{X,x}/\mathfrak{m}_{X,x}^\infty$ is Noetherian for all $x \in X_0$, then we say that X has *Noetherian jet spaces.*

The property of having Noetherian jet spaces is stable under Weil thickenings.

Lemma 2.25. *Let A be a Weil k-superalgebra and X a k-superspace with Noetherian jet spaces. Then X^A has Noetherian jet spaces.*

Proof. Clearly, $\mathfrak{m}_{X^A,x}$ contains $\mathfrak{m}_{X,x} \otimes k$, so $\mathfrak{m}_{X^A,x}^\infty$ contains $\mathfrak{m}_{X,x}^\infty \otimes k$, so there is a surjective homomorphism $\mathcal{O}_{X,x}/\mathfrak{m}_{X,x}^\infty \otimes A \to \mathcal{O}_{X^A,x}/\mathfrak{m}_{X^A,x}^\infty$. The left hand side is finitely generated over the Noetherian ring $\mathcal{O}_{X,x}/\mathfrak{m}_{X,x}^\infty$, so is Noetherian itself. □

Definition 2.26 (Tidiness)**.** Let $j : Y \to X$ be an embedding of superspaces with vanishing ideal $\mathcal{I}$. We say that j is *tidy* if for any $x \in j_0(Y_0)$, any open neighbourhood $U \subseteq X_0$ of x, and any $f \in \mathcal{O}_X(U)$, the following holds:

$$\big(\forall y \in U \cap j_0(Y_0) : f_y \in \mathfrak{m}^\infty_{X,y} + \mathcal{I}_y\big) \;\Rightarrow\; f_x \in \mathcal{I}_x.$$

We say that X is *tidy* if X has Noetherian jet spaces and id_X is a tidy embedding. If X has Noetherian jet spaces, then X is tidy if and only if X has countable girth.

Remark 2.27. Now assume that X has Noetherian jet spaces and let $j : Y \to X$ be an embedding of vanishing ideal $\mathcal{I}$. Then j is tidy if and only if for any $x \in j_0(Y_0)$, any open neighbourhood $U \subseteq X_0$ of x, and any $f \in \mathcal{O}_X(U)$, the following holds:

$$\big(\forall y \in U \cap j_0(Y_0), N \in \mathbb{N} : f_y \in \mathfrak{m}^N_{X,y} + \mathcal{I}_y\big) \;\Rightarrow\; f_x \in \mathcal{I}_x.$$

This follows immediately from Proposition 2.23.

Proposition 2.28 (Tidiness of Weil thickenings)**.** *Let X be a tidy k-superspace and A a Weil k-superalgebra. Then X^A is tidy.*

Proof. By Proposition 2.22, the Weil thickening X^A has countable girth, and by Lemma 2.25, it has Noetherian jet spaces. □

By definition, if $\mathrm{id}_X : X \to X$ is tidy, then X has countable girth. At least if X has Noetherian jet spaces, the same conclusion holds for all tidy embeddings.

Proposition 2.29 (Tidiness of subspaces)**.** *Let X be a superspace and $j : Y \to X$ a tidy embedding. If X has Noetherian jet spaces, then Y is tidy.*

In the *proof*, we first notice the following.

Lemma 2.30. *If $j : Y \to X$ is a not necessarily tidy embedding of superspaces and X has Noetherian jet spaces, then so has Y.*

Proof. Observe simply that for $x = j_0(y)$, the surjection $\mathcal{O}_{X,x} \to \mathcal{O}_{Y,y}/\mathfrak{m}^\infty_{Y,y}$ induced by $j^\sharp$ factors through the Noetherian superring $\mathcal{O}_{X,x}/\mathfrak{m}^\infty_{X,x}$. □

Proof of Proposition 2.29. Let $f \in \mathcal{N}^\infty_Y(V)$, where $V = j_0^{-1}(U)$, $U \subseteq X_0$ being open. Passing to an open cover, we may assume that $f = j^\sharp(g)$ for some $g \in \mathcal{O}_X(U)$.

Let $y \in V$ and $x \coloneqq j_0(y)$. Since $j^\sharp$ is a surjective sheaf map, $\mathfrak{m}_{Y,y} = j^\sharp(\mathfrak{m}_{X,x})$. For any N, we have $f_y \in \mathfrak{m}^N_{Y,y}$. Hence, there are finitely many $g_{kn} \in \mathfrak{m}_{X,x}$, $k = 1, \dots, N$, such that $f_y = j^\sharp\big(\sum_n g_{1n} \cdots g_{Nn}\big)$. So for $h \coloneqq g_x - \sum_n g_{1n} \cdots g_{Nn}$, we have $j^\sharp(h) = 0$, *i.e.* $h \in \mathcal{I}_x$. Thus, $g_x \in \bigcap_{N=1}^\infty (\mathfrak{m}^N_{X,x} + \mathcal{I}_x)$. Since x was arbitrary, it follows that $g_x \in \mathcal{I}_x$ for all $x \in U \cap j_0(Y_0)$. Hence, $f = j^\sharp(g) = 0$. Therefore, Y has countable girth. To see that it has Noetherian jet spaces, we invoke Lemma 2.30. □

Corollary 2.31. *Let X have Noetherian jet spaces and $j : Y \to X$ be an open embedding. Then j is tidy if and only if $j(Y)$ is tidy.*

Proposition 2.32 (Preservation of tidiness)**.** *Let $j : Z \to Y$ and $i : Y \to X$ be embeddings, and Y have Noetherian jet spaces. If j is tidy, then so is $i \circ j$.*

Proof. Denote the vanishing ideals of i, j and $k \coloneqq i \circ j$ by $\mathcal{I}$, $\mathcal{J}$ and $\mathcal{K}$, respectively. Let $U \subset X_0$ be open, $V \coloneqq i_0^{-1}(U)$, and $W \coloneqq j_0^{-1}(V)$. Consider $f \in \mathcal{O}_X(U)$ and $z \in W$. We put $y \coloneqq j_0(z)$ and $x \coloneqq i_0(y)$.

We assume that for all $u \coloneqq i_0(v)$, $v \coloneqq j_0(w)$, $w \in W$, and all $N \in \mathbb{N}$, we have $f_u \in \mathfrak{m}^N_{X,u} + \mathcal{K}_u$. Then there is $g \in \mathfrak{m}^N_{X,u}$ such that $j^\sharp(i^\sharp(f_u - g)) = k^\sharp(f_u - g) = 0$, so that $i^\sharp(f)_v \in i^\sharp(g) + \ker j^\sharp_v \subseteq \mathfrak{m}^N_{Y,v} + \mathcal{J}_v$. Because Y has Noetherian jet spaces by Proposition 2.29 and v, N were arbitrary, we find $i^\sharp(f)_y \in \mathcal{J}_y$, thanks to the tidyness of i. This gives $k^\sharp(f_x) = j^\sharp(i^\sharp(f)_y) = 0$, *i.e.* $f_x \in \mathcal{K}_x$, so that k is tidy. □

Corollary 2.33. *Any embedding of a tidy superspace is tidy.*

Proof. Apply Proposition 2.32 with $Z = Y$ and $j = \mathrm{id}_Y$. □

Proposition 2.34 (Existence of tidying)**.** *Let X be a superspace with Noetherian jet spaces. There is a closed embedding $t : X^\circ \to X$ such that X° is tidy, satisfying the following universal property: Whenever $\varphi : Y \to X$ is a morphism, where Y has countable girth, there is a unique morphism $\varphi^\circ : Y \to X^\circ$ such that $\varphi = t \circ \varphi^\circ$.*

Proof. Define the superspace $X^\circ \coloneqq (X_0, \mathcal{O}_{X^\circ})$ by $\mathcal{O}_{X^\circ} \coloneqq \mathcal{O}_X/\mathcal{N}_X^\infty$. The definition of the embedding $t : X^\circ \to X$ is obvious. Clearly, t is tidy, so that X° is tidy, by Proposition 2.29.

Let $\varphi : Y \to X$ be a morphism, where Y has countable girth. Let $V \subseteq Y_0$ be open, and f a local section of $\mathcal{O}_X$ defined on a neighbourhood U of $\varphi_0(V)$. Assume that for any $y \in V$, $x \coloneqq \varphi_0(y)$, we have $f_x \in \mathfrak{m}_{X,x}^\infty$. Then $\varphi^\sharp(f)_x = \varphi^\sharp(f_x) \in \mathfrak{m}_{Y,y}^\infty$. Because y was arbitrary and Y has countable girth, $\varphi^\sharp(f) = 0$, and φ factors uniquely through t to a morphism $\varphi^\circ : Y \to X^\circ$. □

Definition 2.35 (Tidying)**.** The morphism $t : X^\circ \to X$ and the space X° constructed in Proposition 2.34 are called the *tidying* of X. They are unique up to canonical isomorphism.

Lemma 2.36. *Let X be a superspace with Noetherian jet spaces and $U \subseteq X$ be an open subspace. Then U° is an open subspace of X°.*

Proof. This is obvious by construction. Indeed, $U_0 = (U^\circ)_0$ is open in $X_0 = (X^\circ)_0$ and $\mathcal{O}_{U^\circ} = \mathcal{O}_X|_{U_0}/\mathcal{N}_X^\infty|_{U_0} = (\mathcal{O}_X/\mathcal{N}_X^\infty)|_{U_0} = \mathcal{O}_{X^\circ}|_{U_0}$. □

Proposition 2.37 (Equalisers of tidy superspaces)**.** *Equalisers exist in the category of locally Hausdorff tidy superspaces, and they are tidy embeddings.*

Proof. Let $\phi, \psi : X \to Y$ be morphisms where Y_0 is locally Hausdorff and X has Noetherian jet spaces. Form $Z_0 \coloneqq \{x \in X_0 \mid \phi_0(x) = \psi_0(x)\}$. This is a locally closed subset of X_0 because the diagonal of Y_0 is locally closed. We let $j_0 : Z_0 \to X_0$ be the embedding and set $\mathcal{O}_Z \coloneqq j_0^{-1}\mathcal{O}_X|_U/\mathcal{I}$ where $U \subseteq X_0$ is largest open subset in which Z_0 is closed and $\mathcal{I}$ is the ideal generated by $\mathrm{im}(\phi^\sharp - \psi^\sharp)$. Then $Z \coloneqq (Z_0, \mathcal{O}_Z)$ is a superspace and there is a canonical embedding $j : Z \to X$ such that $\phi \circ j = \psi \circ j$. Since X has Noetherian jet spaces, so does Z, by Lemma 2.30.

Let $t : Z^\circ \to Z$ be the tidying of Z, so that Z° is tidy, and $\jmath^\circ \coloneqq j \circ t$ is a tidy embedding, by Corollary 2.33. Clearly, $\phi \circ \jmath^\circ = \psi \circ \jmath^\circ$. Moreover, if X_0 is locally Hausdorff, then so is Z_0, since j_0 is an embedding.

Now, let W have countable girth and $\varphi : W \to X$ be a morphism such that $\phi \circ \varphi = \psi \circ \varphi$. Then $\varphi^\sharp(\mathcal{I}) = 0$, so that φ factors uniquely through j. Since W has countable girth, φ even factors uniquely through $\jmath^\circ$, by virtue of Proposition 2.34, to a morphism $\varphi^\circ : W \to Z^\circ$. Let $\varrho : W \to Z^\circ$ be another morphism with $\jmath^\circ \circ \varrho = \varphi = \jmath^\circ \circ \varphi^\circ$. Seeing that $\jmath^\circ$ is an embedding and so a monomorphism of superspaces, we find that $\varrho = \varphi^\circ$. □

Proposition 2.38 (Equalisers of open subspaces)**.** *Let $\phi, \psi : X \to Y$ be morphisms of locally Hausdorff tidy superspaces with equaliser $\varphi : Z \to X$. Let $U \subseteq X$ and $V \subseteq Y$ be open subspaces with $U \subseteq \phi^{-1}(V) \cap \psi^{-1}(V)$. Then the equaliser of $\phi|_U, \psi|_U : U \to V$ is an open subspace of Z.*

Proof. Let $W \coloneqq \varphi^{-1}(U)$ and $\varphi|_W : W \to U$ the morphism induced by φ. In view of Corollary 2.31, W is tidy, and it is certainly locally Hausdorff. We claim that $\varphi|_W : W \to U$ is the equaliser of $\phi|_U$ and $\psi|_U$.

Let $\varrho : S \to U$ be a morphism where W is locally Hausdorff and tidy and $\phi|_U \circ \varrho = \psi|_U \circ \varrho$. Suppose that $\sigma \coloneqq j_U \circ \varrho$ factors uniquely through $\varphi : Z \to X$

to a morphism $\sigma' : S \to Z$. Then $\varphi_0(\sigma'_0(S_0)) = \sigma(S_0) \subseteq U_0$, so $\sigma'(S_0) \subseteq W_0$, and σ' factors uniquely through $j_W : W \to Z$ to a morphism $\varrho' : S \to W$. Thus, $j_U \circ \varphi|_W \circ \varrho' = \varphi \circ j_W \circ \varrho' = \sigma = j_U \circ \varrho$, so that $\varphi|_W \circ \varrho' = \varrho$. If $\varrho'' : S \to W$ is another such morphism, then $\varrho' = \varrho''$, because $\varphi|_W$ is an embedding. □

3. Morphisms and local coordinates

This section is devoted to the study of the question to which extent morphisms with range in a supermanifold may be given a coordinate description. We adress this matter by introducing the concept of Leites regularity. It depends on parameters, which indicate whether the supermanifolds in question are smooth, real or complex analytic, and whether their functions take real or complex values.

3.1. **Premanifolds.** In this section, we introduce (pre)manifolds in the traditional manner, using atlases. Then we show how to view them as superspaces. Finally, we state the classical Hadamard lemma, which is instrumental in introducing coordinates into the study of superspaces.

Applications show that it is necessary to consider, apart from cases of the real and complex supermanifolds, superspaces which have the mixed structure of a complex sheaf of superfunctions and an underlying real manifold as their body. Therefor, we will from the very beginning work with a pair $(\mathbb{K}, \Bbbk)$ of fields, where $\Bbbk \subseteq \mathbb{K}$ are both either $\mathbb{R}$ or $\mathbb{C}$.

Definition 3.1 ($\Bbbk$-premanifolds)**.** Let $\varpi = \infty, \omega$ and X_0 a topological space. A *$\Bbbk$-atlas of class $\mathcal{C}^\varpi$* is an open cover (U_i) of X_0, together with topological embeddings $\phi_i : U_i \to \Bbbk^{p_i}$ such that for any i, j,

$$\phi_{ij} \coloneqq \phi_i \circ \phi_j^{-1} : U_{ji} \coloneqq \phi_j(U_i \cap U_j) \to U_{ij}$$

is smooth (for $\varpi = \infty$) or analytic (for $\varpi = \omega$). Two atlases are called *equivalent* if their union is an atlas.

Endowed with an equivalence class of $\Bbbk$-atlases of class $\mathcal{C}^\varpi$, X_0 is called a *$\Bbbk$-premanifold of class $\mathcal{C}^\varpi$*. If in addition, X_0 is Hausdorff, then it is called a *manifold*.

A continuous map $\varphi_0 : X_0 \to Y_0$ of $\Bbbk$-premanifolds of class $\mathcal{C}^\varpi$ is called *of class $\mathcal{C}^\varpi$* (resp. smooth for $\varpi = \infty$, resp. analytic for $\varpi = \omega$), if there exist atlases (U_i, ϕ_i), (V_i, ψ_i) of X_0 resp. Y_0 such that $\varphi_0(U_i) \subseteq V_i$ and $\psi_i \circ \varphi_0 \circ \phi_i^{-1}$ is smooth (for $\varpi = \infty$) resp. analytic (for $\varpi = \omega$).

A map $\varphi_0 : X_0 \to Y_0$ of class $\mathcal{C}^\varpi$ is called an *open embedding* if φ_0 induces a homeomorphism $X_0 \to \varphi_0(X_0)$, endowed with the relative topology from Y_0, and in addition, $\varphi_0(X_0)$ is open in Y_0. In this case, φ_0 is an open map; in fact, it will turn that the open embeddings in this sense coincide with those previously defined, *v.* Corollary 4.3.

We view premanifolds as superspaces for which we impose the regularity conditions and the options for the underlying fields mentioned in the introduction. Accordingly we consider five different classes of premanifolds.

Definition 3.2 (Premanifolds over $(\mathbb{K}, \Bbbk)$)**.** Let $\varpi = \infty, \omega$ and X_0 a $\Bbbk$-premanifold of class $\mathcal{C}^\varpi$. We let $\mathcal{O}_X$ be the sheaf of commutative $\mathbb{K}$-algebras of $\mathbb{K}$-valued functions of class $\mathcal{C}^\varpi$ on X_0. The five possibilities are summarised in the following table:

type of manifold	$\mathbb{K}$	$\Bbbk$	ϖ	$\mathcal{O}_X$
real smooth or $\mathcal{C}^\infty$	$\mathbb{R}$	$\mathbb{R}$	∞	$\mathbb{R}$-valued smooth functions
real analytic or $\mathcal{C}^\omega$	$\mathbb{R}$	$\mathbb{R}$	ω	$\mathbb{R}$-valued real analytic functions
c	$\mathbb{C}$	$\mathbb{R}$	∞	$\mathbb{C}$-valued smooth functions
c analytic	$\mathbb{C}$	$\mathbb{R}$	ω	$\mathbb{C}$-valued real analytic functions
complex analytic	$\mathbb{C}$	$\mathbb{C}$	ω	complex analytic functions

The pair $X \coloneqq (X_0, \mathcal{O}_X)$ will be called a *(pre)manifold over* $(\mathbb{K}, \Bbbk)$ of class $\mathcal{C}^\varpi$. In the case $\varpi = \infty$ resp. $\varpi = \omega$, we say that X is smooth resp. analytic. We also say that X is a (pre)manifold of the type specified in the table. For instance, if $(\mathbb{K}, \Bbbk) = (\mathbb{C}, \mathbb{R})$ and $\varpi = \omega$, we say that X is c *analytic*. Accordingly, the c (pre)manifolds are ordinary smooth (pre)manifolds, endowed with the sheaf of complex-valued smooth functions.

The following lemma, which is immediate from the definitions, shows that the premanifolds from Definition 3.2 are superspaces.

Lemma 3.3. *Let X be a premanifold over $(\mathbb{K}, \Bbbk)$ and $x \in X_0$. Then $\mathcal{O}_{X,x}$ is local and the maximal ideal $\mathfrak{m}_{X,x}$ consists of the germs of functions vanishing at x. Hence, X is a reduced $\mathbb{K}$-rational $\mathbb{K}$-superspace of girth 0; the morphisms of super-ringed spaces associated with maps of class $\mathcal{C}^\varpi$ are morphisms of $\mathbb{K}$-superspaces.*

The $\mathbb{K}$-superspace $X = (X_0, \mathcal{O}_X)$ is called the *$\mathbb{K}$-superspace associated with X_0*. Any such $\mathbb{K}$-superspace will be called a *premanifold over* $(\mathbb{K}, \Bbbk)$ (of class $\mathcal{C}^\varpi$), and a *manifold over* $(\mathbb{K}, \Bbbk)$ (of class $\mathcal{C}^\varpi$) if X_0 is a manifold (*i.e.* Hausdorff).

With any map $\varphi_0 : X_0 \to Y_0$ of $\mathcal{C}^\varpi$ of $\Bbbk$-premanifolds of class $\mathcal{C}^\varpi$, we may associate a morphism $\varphi = (\varphi_0, \varphi^\sharp) : X \to Y$ of the associated $\mathbb{K}$-superspaces, by $\varphi^\sharp(f) \coloneqq f \circ \varphi_0$ for all $f \in \mathcal{O}_Y(U)$, $U \subseteq Y_0$ open. This obviously defines a functor.

Finally, we turn to the Hadamard lemma. It is valid in each of our five categories of premanifolds. To have an efficient formulation, we simply write $\mathbb{A}^p$ for the space $\Bbbk^p$ together with the sheaf $\mathcal{O}_{\mathbb{A}^p}$ of $\mathbb{K}$-valued functions of class $\mathcal{C}^\varpi$, and view it as a manifold over $(\mathbb{K}, \Bbbk)$ of class $\mathcal{C}^\varpi$.

Recall from Definition 2.11 that for any $\mathbb{K}$-superspace X the sheaf of $\Bbbk$-valued superfunctions on X is denoted by $\mathcal{O}_{X,\Bbbk}$.

Proposition 3.4 (Hadamard lemma)**.** *Let $U \subseteq \mathbb{A}^p$ be open. Consider functions $f_1, \dots, f_n \in \mathcal{O}_{\mathbb{A}^p,\Bbbk}(U)$, and set $Y \coloneqq \bigcap_{j=1}^n f_j^{-1}(0)$. Assume that the derivatives $df_1(x) \dots, df_n(x)$ are linearly independent at any $x \in Y$. Then any point of Y possesses an open neighbourhood $V \subseteq U$ such that every $g \in \mathcal{O}_{\mathbb{A}^p}(V)$, $g(V \cap Y) = 0$, admits a representation $g = \sum_{j=1}^n f_j g_j$ for some $g_j \in \mathcal{O}_{\mathbb{A}^p}(V)$. Moreover, we may choose V such that there are $\mathcal{C}^\varpi$ functions $f_{n+1}, \dots, f_p$ on V, for which $f = (f_1, \dots, f_p)$ is a $\mathcal{C}^\varpi$ diffeomorphism onto its open image.*

The following three corollaries of the Hadamard lemma describe the algebraic nature of the sheaf $\mathcal{O}_{\mathbb{A}^p}$.

Corollary 3.5. *Assume the conditions of Proposition 3.4. Take $x \in Y$, and fix $f_{n+1}, \dots, f_p$ as in the proposition. There is an open neighbourhood $V \subseteq U$ of x such that for every $g \in \mathcal{O}_{\mathbb{A}^p}(V)$ and any $N \geqslant 0$, there is a unique polynomial $P = \sum_\alpha P_\alpha(f_{n+1}, \dots, f_p) \cdot T^\alpha$ of degree $\leqslant N$ in $T_1, \dots, T_n$, with coefficients of class $\mathcal{C}^\varpi$, such that $g - P(f_1, \dots, f_n) \in I_Y(V)^{N+1}$, where*

$$I_Y(V) \coloneqq \big\{ h \in \mathcal{O}_{\mathbb{A}^p}(V) \;\big|\; h(V \cap Y) = 0 \big\}.$$

The following two special cases will be of particular importance.

Corollary 3.6 (Pointwise Hadamard lemma)**.** *Let $x \in \mathbb{A}^p_0$ and $f \in \mathcal{O}_{\mathbb{A}^p,x}$. For any $N \geqslant 0$, there is a unique $P \in \mathbb{K}[T_1, \dots, T_p]$, $\deg P \leqslant N$, such that*

$$f - P(t - x) = f - P(t_1 - x_1, \dots, t_p - x_p) \in \mathfrak{m}^{N+1}_{\mathbb{A}^p,x},$$

where $t = (t_1, \dots, t_p)$ is the identity of $\mathbb{A}^p_0$.

Corollary 3.7 (Diagonal Hadamard lemma)**.** *Let $\Delta : \mathbb{A}^p \to \mathbb{A}^{2p}$ be the morphism associated with the diagonal map. Then Δ is a closed embedding and its ideal $\mathcal{I} \subseteq \mathcal{O}_{\mathbb{A}^{2p}}$ is generated by the sections $x_j - y_j$, $j = 1, \dots, p$.*

3.2. **Leites regularity.** In this subsection, we introduce affine superspace, which is the local model for supermanifolds. Affine superspace carries a canonical system of 'coordinate' functions. A natural question is to which extent morphisms to affine superspace are determined by their action on these coordinate functions. This leads to the concept of Leites regularity of superspaces. As we shall see in the following, this gives rise to a remarkably robust and versatile category.

Definition 3.8 (Standard affine superspace)**.** In what follows, we fix the fields $\mathbb{K}$, $\Bbbk$ and $\varpi = \infty, \omega$. We will write $\mathbb{A}^{p|q}$ for the $\mathbb{K}$-superspace $\mathbb{A}^p \times \operatorname{Spec} \bigwedge(\mathbb{K}^q)^*$, where $\mathbb{A}^p$ is the $\mathbb{K}$-superspace associated with the $\Bbbk$-manifold $\Bbbk^p$ of class $\mathcal{C}^\varpi$, and the Weil superalgebra $\bigwedge(\mathbb{K}^q)^* = \mathbb{K}[\theta_1, \dots, \theta_q]$ is the supercommutative $\mathbb{K}$-superalgebra freely generated on q odd generators. We call this the *affine superspace* of dimension $p|q$ (of class $\mathcal{C}^\varpi$ over $(\mathbb{K}, \Bbbk)$).

Note that $\mathbb{A}^{p|0}$ coincides with $\mathbb{A}^p$, as introduced in the previous subsection.

Notation 3.9 (Standard coordinate functions)**.** We denote the standard coordinate functions on $\mathbb{A}^p$ by $t_1, \dots, t_p$. Furthermore, we let $\theta_1, \dots, \theta_q$ be the standard generators of $\bigwedge(\mathbb{K}^q)^* = \mathbb{K}[\theta_1, \dots, \theta_q]$, considered as sections of $\mathcal{O}_{\mathbb{A}^{0|q}}$. We use the same letters if t_i, θ_j are considered as sections of $\mathcal{O}_{\mathbb{A}^{p|q}}$.

We abbreviate the tuple $(t_1, \dots, t_p, \theta_1, \dots, \theta_q)$ by (t, θ). Sometimes, we will not wish to distinguish explicitly between even and odd members of (t, θ). In this case, we will write the standard coordinates $t = (t_a)$ where $a = 1, \dots, p + q$, and we will not impose any particular order on the even and odd members of t. We will say that $t = (t_a)$ is in *standard order* if $|x_a| = \bar{0}$ for all $a \leqslant p$ and $|x_a| = \bar{1}$ for all $a > p$.

On any open $U \subseteq \mathbb{A}^{p|q}_0 = \mathbb{A}^p_0$, every $g \in \mathcal{O}_{\mathbb{A}^{p|q}}(U)$ has a unique expression

$$g = \sum\nolimits_I g_I \theta^I \quad \text{where} \quad \theta^I \coloneqq \theta_{i_1} \cdots \theta_{i_k} \tag{3.1}$$

for $I = (1 \leqslant i_1 < \dots < i_k \leqslant q)$ and $g_I \in \mathcal{O}_{\mathbb{A}^p}(U)$. In particular, the θ_j allow us to consider $\mathcal{O}_{\mathbb{A}^p} \subseteq \mathcal{O}_{\mathbb{A}^{p|q}}$ in a distinguished fashion.

The idea behind the Leites regularity of a $\mathbb{K}$-superspace is to guarantee that a morphism to a standard affine superspace is completely determined by the pullbacks of the standard coordinates under the morphism. It turns out that it is sufficient to demand this for the even standard coordinates.

In what follows, for given $\mathbb{K}$-superspaces X and S, we write

$$X(S) \coloneqq \operatorname{Hom}(S, X)$$

for the set of morphisms $S \to X$ in the category $\mathbf{SSp}_\mathbb{K}$.

Definition 3.10 (Leites regularity and subregularity)**.** We say that a $\mathbb{K}$-superspace X is *(Leites) regular* (of class $\mathcal{C}^\varpi$ over $(\mathbb{K}, \Bbbk)$) if for any open subspace $U \subseteq X$, and any p, the map

$$\mathbb{A}^p(U) = \operatorname{Hom}(U, \mathbb{A}^p) \to \Gamma(\mathcal{O}^p_{U,\Bbbk,\bar{0}}) : \varphi \mapsto (\varphi^\sharp(t_1), \dots, \varphi^\sharp(t_p)), \tag{3.2}$$

where t_j are the coordinate functions on $\mathbb{A}^p$, is bijective. If these maps are all merely injective, then we say that X is *(Leites) subregular*.

We denote the full subcategory of $\mathbf{SSp}_{\mathbb{K}}$ whose objects are the regular superspaces of class $\mathcal{C}^\varpi$ over $(\mathbb{K}, \Bbbk)$ by $\mathbf{SSp}^\varpi_{\mathbb{K},\Bbbk}$. Notice that $\mathbb{A}^p$ is not the p-fold product of $\mathbb{A}^1$ in $\mathbf{SSp}_{\mathbb{K}}$, so that we are obliged to work with $\mathbb{A}^p$. It will turn out that $\mathbb{A}^p$ *is* the p-fold product of $\mathbb{A}^1$ in the subcategory of Leites regular superspaces.

The following lemma shows that for Leites regular superspaces, morphisms to affine superspace are indeed completely described by their tuples of 'component' superfunctions. In fact, it shows that the category of Leites regular superspaces is the largest subcategory of $\mathbf{SSp}_{\mathbb{K}}$ in which Leites's morphism theorem holds: Morphisms to $\mathbb{A}^{p|q}$ are in bijection with tuples of 'component' superfunctions.

Lemma 3.11. *Let X be a Leites regular superspace. The following natural map is a bijection, for any non-negative integers p and q,*

$$\mathbb{A}^{p|q}(X) \to \Gamma(\mathcal{O}_{X,\Bbbk,\bar{0}})^p \times \Gamma(\mathcal{O}_{X,\bar{1}})^q$$
$$\varphi \mapsto (\varphi^\sharp(t_1), \dots, \varphi^\sharp(t_p), \varphi^\sharp(\theta_1), \dots, \varphi^\sharp(\theta_q)).$$

Proof. Since $\mathbb{A}^{p|q} = \mathbb{A}^{p|0} \times \mathbb{A}^{0|q}$ and the statement holds for $q = 0$ by definition, it is sufficient to prove it for $p = 0$. Observe that $\Gamma(\mathcal{O}_{\mathbb{A}^{0|q}}) = \mathbb{K}[\theta_1, \dots, \theta_q]$ is the free supercommutative algebra generated by the odd indeterminates θ_i. Hence, the underlying map of any morphism $\varphi : X \to \mathbb{A}^{0|q}$ necessarily is the constant map $X_0 \to * = (\mathbb{A}^{0|q})_0$, and $\varphi^\sharp(P) = P(\varphi^\sharp(\theta_1), \dots, \varphi^\sharp(\theta_q))$ for any $P \in \mathbb{K}[\theta_1, \dots, \theta_q]$. □

As we will see, the affine superspace $\mathbb{A}^{p|q}$ is Leites regular. This is not hard to prove, but we derive it from more general statements below (Proposition 4.1 and Corollary 4.14). Once it has been established, Lemma 3.11 shows that in the category of Leites regular $\mathbb{K}$-superspaces, we have $\mathbb{A}^{p+r|q+s} = \mathbb{A}^{p|q} \times \mathbb{A}^{r|s}$.

Definition 3.12 (Charts and coordinate systems)**.** Let X be a superspace. A *local chart* is an open embedding of $\mathbb{K}$-superspaces $\varphi : U \to \mathbb{A}^{p|q}$ where $U \subseteq X$ is an open subspace. We say that φ is *defined on* U. If φ is defined on X, then we call φ a *global chart*. The tuple $p|q$ is called the *(graded) dimension* of φ.

Let φ be a local chart. Then the tuple

$$(x, \xi) \coloneqq (x_1, \dots, x_p, \xi_1, \dots, \xi_q) \coloneqq (\varphi^\sharp(t_1), \dots, \varphi^\sharp(t_p), \varphi^\sharp(\theta_1), \dots, \varphi^\sharp(\theta_q))$$

is called a system of *local coordinates* (defined on U). If φ is a global chart, we say that (x, ξ) is a system of *global coordinates*.

Note that $\mathbb{K}$-superspaces admitting enough charts to cover the underlying topological space are automatically Leites regular. Thus, in view of Lemma 3.11, charts and coordinate systems are in bijection. We may and will also call $p|q$ the *(graded) dimension* of (x, ξ).

Occasionally, we will not wish to distinguish in our notation between even and odd members of a system of local coordinates. In this case, we will instead write $x = (x_a)$ where $x_a \coloneqq \varphi^\sharp(t_a)$, $a = 1, \dots, p + q$. We will say that $x = (x_a)$ is in *standard order* if in its definition, (t_a) is in standard order.

In view of Equation (3.1), given a system of local coordinates (x, ξ) defined on $X|_U$, any $f \in \mathcal{O}_X(U)$ has a unique representation

$$f = \sum\nolimits_I f_I \xi^I \quad \text{where} \quad \xi^I \coloneqq \xi_{i_1} \cdots \xi_{i_k} \tag{3.3}$$

and $f_I = x^\sharp(g_I)$ for some $g_I \in \mathcal{O}_{\mathbb{A}^p}(\varphi_0(U))$ where x denotes also the morphism $X|_U \to \mathbb{A}^p$ defined by $x^\sharp(t_i) = x_i$. Observe that the f_I are *not* local sections of $\mathcal{O}_{X_0}$; rather, they are even local sections of $\mathcal{O}_X$.

Although it is justifiable to write $f_I = g_I(x_1, \dots, x_p)$, we will avoid this notation since it is prone to provoke the misunderstanding that the f_I are 'ordinary functions'.

Example 3.13. The standard coordinate system (t, θ) on $\mathbb{A}^{p|q}$ is a global system of coordinates and so is its restriction to any open subspace.

At this point it would be possible to introduce supermanifolds of class $\mathcal{C}^\varpi$ over $(\mathbb{K}, \Bbbk)$ as superspaces which are covered by open subsets admitting global charts. The reason we postpone this is that at the moment we do not have the tools available to also define relative supermanifolds. This will be possible once we have a good understanding of the category of Leites regular $\mathbb{K}$-superspaces and its full subcategory of locally finitely generated $\mathbb{K}$-superspaces (see Section 4.3).

As the following proposition shows, in the framework of Leites regular superspaces, morphisms admit a simple description in terms of coordinates.

Proposition 3.14 (Morphisms *vs.* coordinates)**.** *Let X and Y be Leites regular $\mathbb{K}$-superspaces of class $\mathcal{C}^\varpi$ over $(\mathbb{K}, \Bbbk)$, $y = (y_a)$ a global coordinate system in standard order on Y, and $p|q$ the graded dimension of Y. Then the map*

$$Y(X) \to \big\{x \in \Gamma(\mathcal{O}_X)_{\bar 0}^p \times \Gamma(\mathcal{O}_X)_{\bar 1}^q \;\big|\; x(X_0) \subseteq y(Y_0)\big\} \colon \varphi \mapsto \varphi^\sharp(y)$$

is a bijection. Here, $x(X_0)$—and similarly, $y(Y_0)$—stands for the set

$$x(X_0) \coloneqq \big\{(x_1(o), \dots, x_p(o)) \;\big|\; o \in X_0\big\} \subseteq \mathbb{A}_0^p.$$

Proof. The coordinate system is given by $y = \psi^\sharp(t)$ for a unique global chart $\psi : Y \to \mathbb{A}^{p|q}$, and ψ is an open embedding. Let $x \in \Gamma(\mathcal{O}_X)_{\bar 0}^p \times \Gamma(\mathcal{O}_X)_{\bar 1}^q$ such that $x(X_0) \subseteq y(Y_0)$. Since Y is Leites regular, there exists by Lemma 3.11 a unique morphism $\varrho : X \to \mathbb{A}^{p|q}$ such that $x = \varrho^\sharp(t)$.

The condition $x(X_0) \subseteq y(Y_0)$ guarantees that ϱ factors uniquely through ψ to a morphism $\sigma : X \to Y$, by Proposition 1.12. In particular, $\sigma^\sharp(y) = \varrho^\sharp(t) = x$. The proof of the converse statement is similar. □

Observe that the sets $x(X_0)$ and $y(Y_0)$ in the statement of Proposition 3.14 are contained in $\mathbb{A}_0^p = \Bbbk^p$ (and not only in $\mathbb{K}^p$).

Definition 3.15 (Mapping condition)**.** Let X and Y be regular $\mathbb{K}$-superspaces. Given $x = (x_a) \in \Gamma(\mathcal{O}_X)^{p+q}$ and a global coordinate system $y = (y_a)$ of Y such that $|x_a| = |y_a|$ for all a, the condition

$$x(X_0) \subseteq y(Y_0) \tag{3.4}$$

from Proposition 3.14 is called the *mapping condition* on x and y.

Proposition 3.14 admits an obvious modification for coordinate systems not in standard order. We will use it in this more general form without further notice.

3.3. **Morphisms with non-standard affine target.** Sometimes, it is useful to study the affine superspaces $\mathbb{A}^{p|q}$ independent of coordinates. In this subsection, we digress briefly to show how this can be accomplished in full generality. The basis for this is given by the following definition.

Definition 3.16 (Super-vector spaces over $(\mathbb{K}, \Bbbk)$)**.** Let $V = V_{\bar 0} \oplus V_{\bar 1}$ be a finite-dimensional super-vector space over $\Bbbk$, together with a fixed $\mathbb{K}$-structure on $V_{\bar 1}$. We call V a $(\mathbb{K}, \Bbbk)$-*super-vector space*. For super-vector spaces V, W over $(\mathbb{K}, \Bbbk)$, we set

$$\begin{aligned}\underline{\mathrm{Hom}}_{\mathbb{K},\Bbbk}(V, W) \coloneqq{}& (\mathrm{Hom}_\Bbbk(V_{\bar 0}, W_{\bar 0}) \oplus \mathrm{Hom}_\mathbb{K}(V_{\bar 1}, W_{\bar 1})) \\ &\oplus (\mathrm{Hom}_\Bbbk(V_{\bar 0}, W_{\bar 1}) \oplus \mathrm{Hom}_\Bbbk(V_{\bar 1}, W_{\bar 0})),\end{aligned}$$

with the obvious $\Bbbk$-structure on the even part and the $\mathbb{K}$-structure on the odd part obtained by acting on $W_{\bar{1}}$ in the range and on $V_{\bar{1}}$ in the argument.

In particular, we obtain the *dual super-vector space over* $(\mathbb{K},\Bbbk)$

$$V^* := \underline{\mathrm{Hom}}_{\mathbb{K},\Bbbk}(V,\Bbbk) = \mathrm{Hom}_{\Bbbk}(V_{\bar{0}},\Bbbk)\oplus \mathrm{Hom}_{\mathbb{K}}(V_{\bar{1}},\mathbb{K}).$$

Here, we use the $\mathbb{K}$-structures on $V_{\bar{1}}$ and $\mathrm{Hom}_{\Bbbk}(V_{\bar{1}},\Bbbk)$ to identify the latter canonically with $\mathrm{Hom}_{\mathbb{K}}(V_{\bar{1}},\mathbb{K})$. The even part of $\underline{\mathrm{Hom}}_{\mathbb{K},\Bbbk}(V,W)$ is denoted by $\mathrm{Hom}_{\mathbb{K},\Bbbk}(V,W)$, and its elements are called *morphisms of* $(\mathbb{K},\Bbbk)$*-super-vector spaces*. We also set

$$V\otimes W := (V_{\bar{0}}\otimes_{\Bbbk} W_{\bar{0}}\oplus V_{\bar{1}}\otimes_{\mathbb{K}} W_{\bar{1}})\oplus(V_{\bar{0}}\otimes_{\Bbbk} W_{\bar{1}}\oplus V_{\bar{1}}\otimes_{\Bbbk} W_{\bar{0}}),$$

with the obvious $\Bbbk$- resp. $\mathbb{K}$-structure on the even resp. odd part.

Lemma 3.17. *With the above definitions, finite-dimensional super-vector spaces over* $(\mathbb{K},\Bbbk)$ *form a symmetric monoidal closed category.*

Proof. Let U,V,W be finite-dimensional super-vector spaces over $(\mathbb{K},\Bbbk)$. We have the equations

$$\begin{aligned}
\mathrm{Hom}_{\Bbbk}(U_{\bar{0}}\otimes_{\Bbbk} V_{\bar{0}},W_{\bar{0}}) &= \mathrm{Hom}_{\Bbbk}(U_{\bar{0}},\mathrm{Hom}_{\Bbbk}(V_{\bar{0}},W_{\bar{0}})),\\
\mathrm{Hom}_{\Bbbk}(U_{\bar{1}}\otimes_{\mathbb{K}} V_{\bar{1}},W_{\bar{0}}) &= \mathrm{Hom}_{\mathbb{K}}(U_{\bar{1}},\mathrm{Hom}_{\Bbbk}(V_{\bar{1}},W_{\bar{0}})),\\
\mathrm{Hom}_{\mathbb{K}}(U_{\bar{0}}\otimes_{\Bbbk} V_{\bar{1}},W_{\bar{1}}) &= \mathrm{Hom}_{\Bbbk}(U_{\bar{0}},\mathrm{Hom}_{\mathbb{K}}(V_{\bar{1}},W_{\bar{1}})),\\
\mathrm{Hom}_{\mathbb{K}}(U_{\bar{1}}\otimes_{\Bbbk} V_{\bar{0}},W_{\bar{1}}) &= \mathrm{Hom}_{\mathbb{K}}(U_{\bar{1}},\mathrm{Hom}_{\Bbbk}(V_{\bar{0}},W_{\bar{1}})).
\end{aligned}$$

It follows that there is a natural bijection

$$\mathrm{Hom}_{\mathbb{K},\Bbbk}(U\otimes V,W) = \mathrm{Hom}_{\mathbb{K},\Bbbk}(U,\underline{\mathrm{Hom}}_{\mathbb{K},\Bbbk}(V,W)).$$

From this it follows that the operation $\otimes$ is symmetric and associative. Since $\mathrm{Hom}_{\mathbb{K},\Bbbk}(\Bbbk,U) = U_{\bar{0}}$, it follows that

$$\mathrm{Hom}_{\mathbb{K},\Bbbk}(\Bbbk\otimes U,V) = \mathrm{Hom}_{\mathbb{K},\Bbbk}(\Bbbk,\underline{\mathrm{Hom}}_{\mathbb{K},\Bbbk}(U,V)) = \mathrm{Hom}_{\mathbb{K},\Bbbk}(U,V).$$

Hence, $\Bbbk$ is a tensor unit for $\otimes$, and the assertion follows. □

Construction 3.18 (Affine superspace of a super-vector space)**.** Let V be a $(\mathbb{K},\Bbbk)$-super vector space. The vector space $V_{\bar{0}}$ may be considered as a $\Bbbk$-premanifold. We will denote the associated premanifold over $(\mathbb{K},\Bbbk)$ by the same letter. Let V be the $\mathbb{K}$-superspace $V_{\bar{0}}\times \mathrm{Spec}\bigwedge(V_{\bar{1}})^*$. We call this the *affine superspace* of V. So the reduced space V_0 of V is the premanifold over $(\mathbb{K},\Bbbk)$ given by $V_{\bar{0}}$. The dual V^* is contained in $\Gamma(\mathcal{O}_{V,\Bbbk})$.

To state promised coordinate-free descriptions of morphisms with an affine target cleanly, the following concepts prove useful.

Definition 3.19 ($\mathcal{O}_{X,\Bbbk}$-modules)**.** Let X be a $\mathbb{K}$-superspace. Recall the sheaf $\mathcal{O}_{X,\Bbbk}$ of $\Bbbk$-valued superfunctions, where $\mathcal{O}_{X,\Bbbk}(U)\subseteq\mathcal{O}_X(U)$ is defined by the condition $f(x)\in\Bbbk$ for all $x\in U$, for $U\subseteq X_0$. It is a sheaf of algebras in the symmetric monoidal category of super-vector spaces over $(\mathbb{K},\Bbbk)$.

An $\mathcal{O}_{X,\Bbbk}$*-module* is a sheaf $\mathcal{M}$ with values in the category of super-vector spaces over $(\mathbb{K},\Bbbk)$, which is a graded $\mathcal{O}_{X,\Bbbk}$-module. One has an obvious notion of *morphism of* $\mathcal{O}_{X,\Bbbk}$*-modules* $\mathcal{M}\to\mathcal{N}$. The set of all these will be denoted by $\mathrm{Hom}_{\mathcal{O}_{X,\Bbbk}}(\mathcal{M},\mathcal{N})$. If $\varphi: Y\to X$ is a morphism of $\mathbb{K}$-superspaces and $\mathcal{M}$ is an $(\mathcal{O}_{X,\Bbbk})$-module, then

$$\varphi^*\mathcal{M} := \mathcal{O}_{Y,\Bbbk}\otimes_{\varphi_0^{-1}\mathcal{O}_{X,\Bbbk}}\varphi_0^{-1}\mathcal{M}$$

naturally carries the structure of an $\mathcal{O}_{Y,\Bbbk}$-module.

Construction 3.20 (Free $\mathcal{O}_{X,\Bbbk}$-modules)**.** Let X be a $\mathbb{K}$-superspace and V a finite-dimensional $(\mathbb{K},\Bbbk)$-super-vector space. Let $\mathcal{O}_{X,\Bbbk}\otimes V$ be the tensor product in the category of super-vector spaces over $(\mathbb{K},\Bbbk)$. This is an $\mathcal{O}_{X,\Bbbk}$-module. Similarly, define $\underline{\mathrm{Hom}}_{\mathbb{K},\Bbbk}(V,\mathcal{O}_{X,\Bbbk})$ in the category of super-vector spaces over $(\mathbb{K},\Bbbk)$. Its even part is denoted by $\mathrm{Hom}_{\mathbb{K},\Bbbk}(V,\mathcal{O}_{X,\Bbbk})$.

An $\mathcal{O}_{X,\Bbbk}$-module $\mathcal{P}$ is called *free* if it is isomorphic to $\mathcal{O}_X\otimes V$, for some finite-dimensional $(\mathbb{K},\Bbbk)$-super-vector space V. It is called *locally free* if X_0 admits an open cover, the restrictions of $\mathcal{P}$ to the constituents of which are free.

We have the following canonical isomorphism of $\mathcal{O}_{X,\Bbbk}$-modules,

$$\underline{\mathrm{Hom}}_{\mathbb{K},\Bbbk}(V,\mathcal{O}_{X,\Bbbk})\cong\mathcal{O}_{X,\Bbbk}\otimes V^*.$$

In particular, $\underline{\mathrm{Hom}}_{\mathbb{K},\Bbbk}(V,\mathcal{O}_{X,\Bbbk})$ is a free $\mathcal{O}_{X,\Bbbk}$-module.

With the help of this terminology, we can now give a coordinate-free description of morphisms with affine target.

Corollary 3.21. *Let X be a regular $\mathbb{K}$-superspace and V a $(\mathbb{K},\Bbbk)$-super-vector space. Let V also denote the associated affine superspace. Then there is a bijection*

$$V(X)\longrightarrow\Gamma\big(\mathrm{Hom}_{\mathbb{K},\Bbbk}(V^*,\mathcal{O}_{X,\Bbbk})\big)=\Gamma\big((\mathcal{O}_{X,\Bbbk}\otimes V)_{\bar0}\big)$$

which is natural in X and V. Under this map,

$$V_{\bar0}(X)\cong\Gamma\big(\mathrm{Hom}_{\Bbbk}(V_{\bar0}^*,\mathcal{O}_{X,\Bbbk,\bar0})\big)=\Gamma\big(\mathcal{O}_{X,\Bbbk,\bar0}\otimes_{\Bbbk}V_{\bar0}\big),$$
$$V_{\bar1}(X)\cong\Gamma\big(\mathrm{Hom}_{\mathbb{K}}(V_{\bar1}^*,\mathcal{O}_{X,\bar1})\big)=\Gamma\big(\mathcal{O}_{X,\bar1}\otimes_{\mathbb{K}}V_{\bar1}\big).$$

More generally, for each open subspace $U\subseteq V$, there is a natural bijection

$$\begin{aligned}U(X)\cong&\big\{f\in\Gamma(\mathrm{Hom}_{\mathbb{K},\Bbbk}(V^*,\mathcal{O}_{X,\Bbbk}))\;\big|\;\forall x\in X_0\exists u\in U_0:f(\cdot)(x)=\langle\cdot,u\rangle\big\}\\=&\big\{f=\textstyle\sum_i f_i\otimes v_i\in\Gamma((\mathcal{O}_{X,\Bbbk}\otimes V)_{\bar0})\;\big|\;\forall x\in X_0\,\exists u\in U_0:u=\sum_i f_i(x)v_i\big\}\end{aligned}$$

Here, $\langle\cdot,\cdot\rangle$ denotes the canonical pairing of V^* and V.

4. Singular superspaces and relative supermanifolds

This section contains our main results. Firstly, we show that Leites regularity of superspaces is preserved under colimits and also under the passage to subspaces and thickenings, at least if they are subregular. We then construct the subcategory of locally finitely generated superspaces. These admit finite limits, and are stable both under passage to tidy subspaces and Weil thickenings.

Thus, within the category of locally finitely generated superspaces, we may construct the category of relative supermanifolds over a possibly singular base. This also provides a natural framework for the study of Weil functors, which give a uniform description of such natural geometric objects as the even and odd tangent bundle and higher order versions of these.

4.1. **Constructions of Leites regular superspaces.** So far, we have established that Leites regularity is a desirable property for $\mathbb{K}$-superspaces. However, we have not given any examples of superspaces with this property. In this subsection, we will derive a number of stability results that will furnish a generous supply.

We start by showing that premanifolds are Leites regular. Next, we show that Leites regularity is stable under colimits. Moreover, we investigate to which extent it is preserved under taking subspaces and thickenings, for example, Weil thickenings.

For the sake of brevity, from now on, we will drop the prefix 'Leites' and just speak of 'regular' resp. 'subregular' superspaces.

Proposition 4.1 (Regularity of premanifolds)**.** *Let X be a premanifold over $(\mathbb{K},\Bbbk)$. Then X is regular. In particular, $\mathbb{A}^p$ is regular.*

For the *proof*, we make the following observation, which is of independent interest.

Lemma 4.2. *Any $\mathbb{K}$-superspace of countable girth is subregular.*

Proof. Let X have countable girth and let morphisms $\varphi, \psi : X \to \mathbb{A}^p$ be given such that $\varphi^\sharp(t_j) = f_j = \psi^\sharp(t_j)$. Then

$$t_j(\varphi_0(x)) = j^\sharp_{X_0}\varphi^\sharp(t_j)(x) = j^\sharp_{X_0}\psi^\sharp(t_j)(x) = t_j(\psi_0(x))$$

for any $x \in X_0$ and $j = 1, \dots, p$, so that $\varphi_0 = \psi_0$.

Let f be a local section of $\mathcal{O}_{\mathbb{A}^p}$. Let $N \in \mathbb{N}$ and $x \in X_0$ be arbitrary, and consider the point $y \coloneqq \varphi_0(x) = \psi_0(x)$. By Corollary 3.6, there is a polynomial $P \in \mathbb{K}[t_1 - y_1, \dots, t_p - y_p]$ such that $(f - P)_y \in \mathfrak{m}^{N+1}_{\mathbb{A}^p,y}$. Then

$$\varphi^\sharp(f)_x - \psi^\sharp(f)_x = \varphi^\sharp(f - P)_x - \psi^\sharp(f - P)_x \in \mathfrak{m}^{N+1}_{X,x}.$$

Since x and N were arbitrary, we conclude that $\varphi^\sharp(f) - \psi^\sharp(f)$ are local sections of $\mathcal{N}^\infty_X$, which vanishes by assumption. It follows that $\varphi^\sharp(f) = \psi^\sharp(f)$. □

Proof of Proposition 4.1. Given functions $f_1, \dots, f_p \in \Gamma(\mathcal{O}_{X,\Bbbk})$, there is a unique premanifold map $\varphi_0 : X_0 \to \Bbbk^p$ of class $\mathcal{C}^\varpi$ with components $t_j \circ \varphi_0 = f_j$. The associated morphism $\varphi : X \to \mathbb{A}^p$ (compare the remarks following Lemma 3.3) satisfies $\varphi^\sharp(t_j) = f_j$, proving surjectivity. All open subspaces of X are premanifolds, so Lemma 4.2 implies the claim. □

Corollary 4.3. *Consider the functor which sends a $\Bbbk$-premanifold of class $\mathcal{C}^\varpi$ to its associated $\mathbb{K}$-superspace. It maps open embeddings of premanifolds to open embeddings of $\mathbb{K}$-superspaces, is fully faithful, and its image consists of the premanifolds over $(\mathbb{K}, \Bbbk)$ of class $\mathcal{C}^\varpi$.*

Proof. Let $\varphi_0 : X_0 \to Y_0$ be a manifold map of class $\mathcal{C}^\varpi$ which is an open embedding of topological spaces. Then the associated morphism $\varphi : X \to Y$ is an open embedding of $\mathbb{K}$-superspaces. Hence, the statement follows from Proposition 1.20 and Proposition 4.1. □

The category of regular superspaces is quite robust.

Proposition 4.4 (Colimits of regular superspaces)**.** *The full subcategory of $\mathbf{SSp}_\mathbb{K}$ of regular (resp. subregular) $\mathbb{K}$-superspaces is cocomplete.*

Proof. Let I be a small category, $F : I \to \mathbf{SSp}_\mathbb{K}$ a functor, and $X = \varinjlim_I F$ in $\mathbf{SSp}_\mathbb{K}$ (*v.* Proposition 1.15). Denote the canonical morphisms $F(i) \to X$ by j_i. Assume that F takes values in subregular superspaces. Let $\varphi, \psi : X \to \mathbb{A}^p$ be morphisms such that $f_k \coloneqq \varphi^\sharp(t_k) = \psi^\sharp(t_k)$, $k = 1, \dots, p$. For any i, we form $f^i_k \coloneqq j^\sharp_i(f_k)$. Then $(\varphi \circ j_i)^\sharp(t_k) = f^i_k = (\psi \circ j_i)^\sharp(t_k)$, $k = 1, \dots, p$, so by assumption, we have $\varphi \circ j_i = \psi \circ j_i$. Since i was arbitrary, this shows $\varphi = \psi$, so X is subregular.

Assume now that F even takes values in regular superspaces, fix even superfunctions $f_1, \dots, f_p \in \Gamma(\mathcal{O}_{X,\Bbbk,\bar{0}})$, and form f^i_k as above. By assumption, there are unique $\varphi_i : F(i) \to \mathbb{A}^p$ such that $\varphi^\sharp_i(t_k) = f^i_k$, $k = 1, \dots, p$. For any $\chi : i \to i'$ in I, we have

$$F(\chi)^\sharp(f^{i'}_k) = (j_{i'} \circ F(\chi))^\sharp(t_k) = j^\sharp_i(t_k) = f^i_k$$

By the uniqueness, this implies $\varphi_{i'} \circ F(\chi) = \varphi_i$, so there exists a unique morphism $\varphi : X \to \mathbb{A}^p$ such that $\varphi \circ j_i = \varphi_i$ for all i. Then $j^\sharp_i(\varphi^\sharp(t_k)) = \varphi^\sharp_i(t_k) = f^i_k$ for any i. This implies $\varphi^\sharp(t_k) = f_k$, $k = 1, \dots, p$, in view of Remark 1.16. □

In particular, one may consider infinite versions of the affine superspaces $\mathbb{A}^p$.

Notation 4.5 (Infinite affine superspace $\mathbb{A}^\infty$)**.** For any integers $p' > p \geqslant 0$, let $j_{p'p} : \mathbb{A}^p \to \mathbb{A}^{p'}$ be the morphism induced by the map $x \mapsto (x, 0_{p'-p})$. Then the family $(\mathbb{A}^p, j_{p'p})$ is an inductive system of $\mathbb{K}$-superspaces. Let $\mathbb{A}^\infty$ denote its colimit.

By Proposition 4.4, $\mathbb{A}^\infty$ is a regular superspace, and by Remark 1.16, there is an infinite sequence $t_1, t_2, \ldots \in \Gamma(\mathcal{O}_{\mathbb{A}^\infty,\mathbb{k}})$ of coordinate functions. It is determined by $j_p^\sharp(t_j) = t_j$ for all $j \leqslant p$, where $j_p : \mathbb{A}^p \to \mathbb{A}^\infty$ are the natural morphisms.

The proof of the following lemma is similar to that of Proposition 4.4.

Lemma 4.6. *Let X be a $\mathbb{K}$-superspace. Then X is regular (resp. subregular) if and only if for each open subspace $U \subseteq X$, the following natural map is bijective (resp. injective):*

$$\mathbb{A}^\infty(U) \to \prod_{k=1}^{\infty} \Gamma(\mathcal{O}_{U,\mathbb{k},\bar{0}}) : \varphi \mapsto (\varphi^\sharp(t_1), \varphi^\sharp(t_2), \ldots). \tag{4.1}$$

We now derive an explicit description of the infinitesimal normal neighbourhoods of the diagonal of $\mathbb{A}^p$.

Notation 4.7. Let $p \in \mathbb{N}$ and $\Delta : \mathbb{A}^p \to \mathbb{A}^{2p}$ be the diagonal morphism. Denote by $(\mathbb{A}^p)^{(n)}_\Delta$, for $n \in \mathbb{N} \cup \infty$, the infinitesimal normal neighbourhood of order n of $\mathbb{A}^p$ with respect to Δ.

Let J^p_n denote the Weil algebra $\mathbb{K}[T_1, \ldots, T_p]/(T_1, \ldots, T_p)^{n+1}$ for $n \in \mathbb{N}$, and let $J^p_\infty = \mathbb{K}[\![T_1, \ldots, T_p]\!]$ be the corresponding projective limit. Here, $\mathbb{K}[\![T_1, \ldots, T_p]\!]$ is the $\mathbb{K}$-algebra of formal power series in the algebraically independent indeterminates T_j. We call J^p_n, J^p_∞ *Ehresmann jet algebras*.

Proposition 4.8 (Infinitesimal neighbourhoods of the diagonal)**.** *Let $p, n \in \mathbb{N}$. We have $(\mathbb{A}^p)^{(n)}_\Delta = \mathbb{A}^p \times \operatorname{Spec} J^p_n$, so that $\mathcal{O}_{(\mathbb{A}^p)^{(n)}_\Delta} = \mathcal{O}_{\mathbb{A}^p} \otimes J^p_n$. The latter statement also holds for $n = \infty$, so that $\mathcal{O}_{(\mathbb{A}^p)^{(\infty)}_\Delta} = \mathcal{O}_{\mathbb{A}^p}[\![T_1, \ldots, T_p]\!]$.*

Proof. Let us first assume that n is finite. We let $f_j \coloneqq x_j - y_j$ for $j = 1, \ldots, p$ and $f_j \coloneqq x_{j-p}$ for $j = p+1, \ldots, 2p$. Then the assertion follows immediately from Corollary 3.5. The superspaces $(\mathbb{A}^p)^{(n)}_\Delta$ form an inductive system for fixed p. Since $\mathcal{O}_{\mathbb{A}^p} \otimes J^p_\infty$ is the projective limit of $\mathcal{O}_{\mathbb{A}^p} \otimes J^p_n$, for any p, the assertion follows. □

We now investigate when the regularity of $\mathbb{K}$-superspaces is inherited by subspaces. To that end, consider the following definition.

Definition 4.9 (Local factorisation)**.** Let $f : X \to Y$, $h : X \to Z$ be morphisms of $\mathbb{K}$-superspaces. We say that h *locally factors through* f resp. *locally factors uniquely* if there are open covers X_i, Y_i of X, Y such that $X_i \subseteq f^{-1}(Y_i)$ and $h : X_i \to Z$ factors resp. factors uniquely through $f : X_i \to Y_i$, for any i.

Proposition 4.10 (Regularity of subspaces)**.** *Let $j : Y \to X$ be an embedding of $\mathbb{K}$-superspaces where X is regular. The following are equivalent:*

(i) *The $\mathbb{K}$-superspace Y is regular;*
(ii) *morphisms from open subspaces of Y to $\mathbb{A}^p$ locally factor through j; and*
(iii) *the $\mathbb{K}$-superspace Y is subregular.*

The equivalent conditions hold if $j : Y \to X$ is locally retractable.

The *proof* uses the following lemma, which is based on ideas from the theory of ideals of $\mathcal{C}^\infty$-functions, *v.* [9, Chapter I, Proposition 1.2].

Lemma 4.11. *Let X be regular, $\varphi : Y \to X$ and $\phi, \psi : X \to \mathbb{A}^p$ be morphisms of $\mathbb{K}$-superspaces. If $(\phi \circ \varphi)^\sharp(t_i) = (\psi \circ \varphi)^\sharp(t_i)$, for $i = 1, \ldots, p$, then $\phi \circ \varphi = \psi \circ \varphi$.*

Proof. Since X is regular, we may define $\varrho : X \to \mathbb{A}^{2p}$ by the requirement that $\varrho^\sharp(x_i) = \phi^\sharp(t_i)$ and $\varrho^\sharp(y_i) = \psi^\sharp(t_i)$. Similarly, since $\mathbb{A}^{2p}$ is regular, there are unique morphisms $p_1, p_2 : \mathbb{A}^{2p} \to \mathbb{A}^p$ such that $p_1^\sharp(t_i) = x_i$ and $p_2^\sharp(t_i) = y_i$. Obviously, $p_1 \circ \varrho = \phi$ and $p_2 \circ \varrho = \psi$, where again, we have used the regularity.

Now, let $k \in \mathcal{O}_{\mathbb{A}^p}(W)$ where $W \subseteq \mathbb{A}^p_0$ is open. According to Corollary 3.7, there exist functions k_i defined on some open neighbourhood of $\Delta_0(W) \subseteq \mathbb{A}^p_0 \times \mathbb{A}^p_0$ with

$$p_1^\sharp(k) - p_2^\sharp(k) = \sum\nolimits_{i=1}^{p} (x_i - y_i) \cdot k_i.$$

We thus compute

$$\phi^\sharp(k) - \psi^\sharp(k) = \varrho^\sharp(p_1^\sharp(k) - p_2^\sharp(k)) = \sum\nolimits_{i=1}^{p} (\phi^\sharp(t_i) - \psi^\sharp(t_i))\varrho^\sharp(k_i),$$

so $\varphi^\sharp\phi^\sharp(k) = \varphi^\sharp\psi^\sharp(k)$. Since k was arbitrary, we have $\phi \circ \varphi = \psi \circ \varphi$. □

Proof of Proposition 4.10. We may assume that j is a closed embedding, and by passing to open subspaces, it is sufficient to consider the maps $\mathbb{A}^p(Y) \to \Gamma(\mathcal{O}_{Y,\Bbbk,\bar{0}})^p$. Let $f_1, \dots, f_p \in \Gamma(\mathcal{O}_{Y,\Bbbk,\bar{0}})$. There exist an open cover (U_i) of X_0 and even superfunctions $g_{ij} \in \mathcal{O}_X(U_i)_{\bar{0}}$, $j = 1, \dots, p$, such that $j^\sharp(g_{ij}) = f_j|_{V_i}$, where $V_i \coloneqq j_0^{-1}(U_i)$.

By the assumption on X, there are morphisms $\varphi_i : X|_{U_i} \to \mathbb{A}^p$ such that $\varphi_i^\sharp(t_j) = g_{ij}$. In view of Lemma 4.11, the morphisms $\varphi_k \circ j$ and $\varphi_\ell \circ j$ coincide on $Y|_{V_k \cap V_\ell}$. Hence, by Proposition 1.20, there is a morphism $\varphi : Y \to \mathbb{A}^p$ such that $\varphi = \varphi_i \circ j$ on $Y|_{V_i}$. In particular, we have $\varphi^\sharp(t_j) = f_j$. Hence, the canonical map $\mathbb{A}^p(Y) \to \Gamma(\mathcal{O}_{Y,\Bbbk,\bar{0}})^p$ is surjective for any p, and (i) and (iii) are equivalent.

On the other hand, let $\psi : Y \to \mathbb{A}^p$ be another morphism such that $\psi^\sharp(t_j) = f_j$, and assume that ψ locally factors through j. Possibly after refining the open cover (U_i), $\psi : Y|_{V_i} \to \mathbb{A}^p$ factors through $j : Y|_{V_i} \to X|_{U_i}$ by morphisms $\psi_i : X|_{U_i} \to \mathbb{A}^p$. Then $h_{ij} \coloneqq \psi_i^\sharp(t_j)$ satisfies $j^\sharp(h_{ij}) = f_j|_{V_i}$. Applying Lemma 4.11 and Proposition 1.20 again, we find that $\psi = \varphi$.

Hence, if any morphism from an open subspace of Y to $\mathbb{A}^p$ locally factors through j, then Y is regular. In particular, (ii) implies (i). Conversely, assume that Y is regular, and let $\varphi : Y \to \mathbb{A}^p$ be a morphism. (By the definition of regularity, we may assume that φ is defined on all of Y.) Let $f_j \coloneqq \varphi^\sharp(t_j)$ and (U_i), (V_i), g_{ij} be as above. Define $\varphi_i : X|_{U_i} \to \mathbb{A}^p$ by the condition $\varphi_i^\sharp(t_j) = g_{ij}$. Then the regularity of Y shows that $\varphi = \varphi_i \circ j$ on $Y|_{V_i}$, so that φ locally factors through j.

Finally, if $j : X \to Y$ is locally retractable, then (ii) clearly holds. □

Corollary 4.12. *Let $p, n \in \mathbb{N} \cup \infty$. The $\mathbb{K}$-superspaces $(\mathbb{A}^p)^{(n)}_\Delta$ are regular.*

Proof. In view of Proposition 4.4, and because inductive limits commute, it is sufficient to prove this for p and n finite. The thickening $j^{(0n)} : \mathbb{A}^p \to (\mathbb{A}^p)^{(n)}_\Delta$ has finite girth. The reduction of $(\mathbb{A}^p)^{(n)}_\Delta$ is $\mathbb{A}^p$ itself. Therefore, $(\mathbb{A}^p)^{(n)}_\Delta$ has finite girth, by Proposition 2.2, and so, is subregular, in view of Lemma 4.2. Proposition 4.10 applies to the embedding $(\mathbb{A}^p)^{(n)}_\Delta \to \mathbb{A}^{2p}$ induced by Δ, which implies the claim. □

Here is a partial converse of the previous proposition.

Proposition 4.13 (Regularity of thickenings)**.** *Let X and Y be $\mathbb{K}$-superspaces where X is subregular, and $j : Y \to X$ be a locally retractable thickening of finite girth. If Y is regular, then so is X.*

Proof. In view of Proposition 1.20, the question is local, so that we may assume, by passing to an open subspace, that j admits a retraction $r : X \to Y$ and is of finite girth N. Then $X = Y^{(N)}$, where $Y^{(n)}$ denotes the normal infinitesimal neighbourhood of Y in X of order n.

Let $f_1, \dots, f_p \in \Gamma(\mathcal{O}_{X,\Bbbk,\bar{0}})$ and set $g_j \coloneqq j^\sharp(f_j)$. Denote by $\varphi : Y \to \mathbb{A}^p$ the unique morphism such that $\varphi^\sharp(t_j) = g_j$, and set $\varepsilon_j \coloneqq r^\sharp(g_j) - f_j$. Since $j^\sharp(\varepsilon_j) = 0$,

the ε_j are by assumption nilpotent elements of $\Gamma(\mathcal{O}_X)$. By Proposition 4.8, we may define a morphism $\psi^\infty : X \to (\mathbb{A}^p)^{(\infty)}_\Delta$ by setting $\psi^\infty_0 := \varphi_0$ and for any $h \in \mathcal{O}_{\mathbb{A}^p}(U)$

$$\psi^{\infty\sharp}(h \cdot T^\alpha) := r^\sharp(\varphi^\sharp(h)) \cdot \varepsilon^\alpha,$$

where we use the familiar multi-index notation.

Let $r^n_{\mathbb{A}} : (\mathbb{A}^p)^{(n)}_\Delta \to \mathbb{A}^p$ be the morphism $\Delta^{(n)} : (\mathbb{A}^p)^{(n)}_\Delta \to \mathbb{A}^{2p}$, composed with the projection $p_2 : \mathbb{A}^{2p} \to \mathbb{A}^p$. We define the morphism $\psi := r^\infty_{\mathbb{A}} \circ \psi^\infty : X \to \mathbb{A}^p$. Then $r^{\infty\sharp}_{\mathbb{A}}(t_j) = y_j = x_j - 1 \cdot (x_j - y_j)$. Considered as a section of $\mathcal{O}_{\mathbb{A}^p}[\![T_1, \dots, T_p]\!]$, the latter quantity equals $t_j \cdot T^0 - 1 \cdot T_j$. Hence,

$$\psi^\sharp(t_j) = r^\sharp(g_j) - \varepsilon_j = f_j,$$

so that ψ meets the requirements. Since X is subregular, it is regular. □

Corollary 4.14 (Regularity of Weil thickenings)**.** *Let X be a regular $\mathbb{K}$-superspace of countable girth and A a Weil $\mathbb{K}$-superalgebra. Then the Weil thickened $\mathbb{K}$-superspace X^A is regular. In particular,* Spec A *itself is regular.*

Proof. By Proposition 2.22, X^A also has countable girth. By Lemma 4.2, X^A is subregular. Hence, Proposition 4.13 applies, and the assertion follows. □

4.2. **Locally finitely generated superspaces.** The regular spaces we have considered so far have many desirable permanence properties under the assumption that subregularity is preserved. However, subregularity is not automatically preserved under thickenings and embeddings, and relatedly, (fibre) products do not exist.

We will now single out the full subcategory of locally finitely generated superspaces, which turns out to be better behaved on both accounts: it is stable under Weil thickenings and admits fibre products.

Definition 4.15 (Finitely generated superspaces)**.** Let X be a $\mathbb{K}$-superspace. If there exists a tidy embedding $j : X \to \mathbb{A}^p \times \operatorname{Spec} A$ for some Weil $\mathbb{K}$-superalgebra A, then X is called *finitely generated*. It is called *locally finitely generated* if it admits a cover by finitely generated open subspaces.

Proposition 4.16 (Finite generation implies tidiness)**.** *Let X be a locally finitely generated $\mathbb{K}$-superspace. Then X is tidy and hence regular.*

The *proof* relies on the following fact.

Lemma 4.17. *Let $p \in \mathbb{N}$. The $\mathbb{K}$-superspace $\mathbb{A}^p$ has Noetherian jet spaces.*

Proof. If $\varpi = \omega$, then $\mathcal{O}_{\mathbb{A}^p,x}$ is the convergent power series ring in p indeterminates, so $\mathfrak{m}^\infty_{\mathbb{A}^p,x} = 0$ and $\mathcal{O}_{\mathbb{A}^p,x}$ is itself Noetherian. If $\varpi = \infty$, then

$$\mathcal{O}_{\mathbb{A}^p,x}/\mathfrak{m}^\infty_{\mathbb{A}^p,x} = \mathbb{K}[\![T_1, \dots, T_p]\!]$$

by É. Borel's theorem, and this is again a Noetherian ring. □

Proof of Proposition 4.16. By Lemma 4.17 and Lemma 2.25, any Weil thickening of $\mathbb{A}^p$ has Noetherian jet spaces. By the assumption and Proposition 2.29, X is tidy, since this is a local property. By Lemma 4.2 and Proposition 4.10, X is regular. □

Corollary 4.18 (Locally finite generation of Weil thickenings)**.** *The category of (locally) finitely generated $\mathbb{K}$-superspaces is stable under Weil thickenings.*

Proof. Let $\varphi : X \to \mathbb{A}^p \times \operatorname{Spec} A$ be a tidy embedding and B a Weil $\mathbb{K}$-superalgebra. Applying the functor $(-)^B = (-) \times \operatorname{Spec} B$, we obtain by Proposition 2.20 a morphism $\varphi^B : X^B \to \mathbb{A}^p \times \operatorname{Spec}(A \otimes B)$. Evidently, it is an embedding. By Proposition 4.16, X is tidy, hence, so is X^B, by virtue of Proposition 2.28. Then Corollary 2.33 shows that φ^B is tidy. □

Note that $\mathbb{A}^{p|q}$ is a Weil thickening of $\mathbb{A}^p$, so Corollary 4.18 says in particular that $\mathbb{A}^{p|q}$ is locally finitely generated.

Corollary 4.19 (Finite generation and affine embeddings)**.** *Let X be finitely generated. Then X admits a tidy embedding into some $\mathbb{A}^{p|q}$.*

Proof. By Proposition 4.16, X is tidy, and any embedding into $\mathbb{A}^{p|q}$ will hence be tidy and regular, in view of Corollary 2.33. It suffices to show that any $\mathbb{A}^r \times \operatorname{Spec} A$, where A is a Weil $\mathbb{K}$-superalgebra, can be embedded into some $\mathbb{A}^{p+r|q}$.

Since A is finitely generated by nilpotent elements, there exists a surjective homomorphism from $B \otimes C$ onto A, where $C \coloneqq \mathbb{K}[\Theta_1, \dots, \Theta_q]$ and

$$B \coloneqq \mathbb{K}[T_1, \dots, T_p]/(T_1^{\alpha_1} \cdots T_p^{\alpha_p} | \alpha_1 + \cdots + \alpha_p = N),$$

for some non-negative integers p, q, and N, and even resp. odd indeterminates T_i resp. Θ_j. We may assume $A = B \otimes C$. Since $\operatorname{Spec} C = \mathbb{A}^{0|q}$, we may assume $A = B$.

Define $\varphi : \mathbb{A}^r \times \operatorname{Spec} A \to \mathbb{A}^{p+r}$ by letting $\varphi_0 \coloneqq \mathrm{id}_{\mathbb{k}^r} \times 0$ and $\varphi^\sharp$ be the canonical projection $\mathcal{O}_{\mathbb{A}^{p+r}} \to \mathcal{O}_{\mathbb{A}^{p+r}}/\mathcal{I}^N$ where $\mathcal{I}(U) \coloneqq \mathcal{O}_{\mathbb{A}^{p+r}}(U)$ if $U \cap (\mathbb{k}^r \times 0) = \varnothing$ and

$$\mathcal{I}(U) \coloneqq \left\{ f \in \mathcal{O}_{\mathbb{A}^{p+r}}(U) \;\middle|\; f(\mathbb{k}^r \times 0) = 0 \right\}$$

otherwise. Then by Corollary 3.5, $\varphi_0^{-1}(\mathcal{O}_{\mathbb{A}^{p+r}}/\mathcal{I}^N) \cong \mathcal{O}_{\mathbb{A}^r} \otimes A$. Thus, φ is indeed a well-defined embedding $\mathbb{A}^r \times \operatorname{Spec} A \to \mathbb{A}^{p+r}$. □

Corollary 4.20 (Characterisation of finite generation)**.** *Let X be a $\mathbb{K}$-superspace. The following are equivalent:*

(i) *X is locally finitely generated;*

(ii) *X is tidy and admits an open cover by subspaces embeddable into $\mathbb{A}^p \times \operatorname{Spec} A$, for some integers p and Weil $\mathbb{K}$-superalgebras A;*

(iii) *X is tidy and admits an open cover by subspaces embeddable into $\mathbb{A}^{p|q}$, for some integers p and q.*

Proof. We have (i) ⇒ (ii) by Proposition 4.16 and Corollary 4.19, and (ii) ⇒ (iii) is obvious. Finally, (ii) ⇒ (i) follows from Corollary 2.33 and the definitions. □

Proposition 4.21 (Finite generation of reduction and body)**.** *Let X be locally finite generated. Then so are X_0 and $X_{\bar{0}}$.*

Proof. We may assume that X is finitely generated, so that, taking Corollary 4.19 into consideration, there is a tidy embedding $\varphi : X \to \mathbb{A}^{p|q}$. Then $(\mathbb{A}^{p|q})_0 = (\mathbb{A}^{p|q})_{\bar{0}} = \mathbb{A}^p$ is reduced and in particular even.

Since $\varphi \circ j_{X_{\bar{0}}} = j_{(\mathbb{A}^{p|q})_{\bar{0}}} \circ \varphi_{\bar{0}}$ is an embedding, $\varphi_{\bar{0}} : X_{\bar{0}} \to \mathbb{A}^p$ is an embedding. Similarly, so is $\varphi_0 : X_0 \to \mathbb{A}^p$. In particular, by Lemma 2.30, both X_0 and $X_{\bar{0}}$ have Noetherian jet spaces.

Since X_0 has girth zero, it is finitely generated, by Corollary 4.20. Reasoning similarly, it remains to be shown that $X_{\bar{0}}$ has countable girth, and to that end, it is sufficient that the embedding $\varphi_{\bar{0}}$ be tidy.

So, let $\mathcal{I}$ denote the vanishing ideal of φ and $\bar{\mathcal{I}}$ that of $\varphi_{\bar{0}}$. Fix $f \in \mathcal{O}_{\mathbb{A}^p}(U)$ such that $f_x \in \mathfrak{m}^\infty_{\mathbb{A}^p,x} + \bar{\mathcal{I}}_x$ for all $x \in U \cap \varphi_0(X_0)$. We may consider local sections $\mathcal{O}_{\mathbb{A}^p}$ as local sections of of $\mathcal{O}_{\mathbb{A}^{p|q}}$, as in Notation 3.9. The sheaf map $\mathcal{O}_{\mathbb{A}^p} \to \mathcal{O}_{\mathbb{A}^{p|q}}$ considered there is local on the stalks. Thus, if $f_x = g_x + h_x$ where $g_x \in \mathfrak{m}^\infty_{\mathbb{A}^p,x}$ and $h_x \in \bar{\mathcal{I}}_x$, then $g_x \in \mathfrak{m}^\infty_{\mathbb{A}^{p|q},x}$. Moreover, $\varphi^\sharp(h_x) = \varphi^\sharp_{\bar{0}}(h_x) = 0$, so that $f_x \in \mathfrak{m}^\infty_{\mathbb{A}^{p|q},x} + \mathcal{I}_x$ for all $x \in U \cap \varphi_0(X_0)$. By the assumption on φ, this implies that $\varphi^\sharp(f) = 0$, and in particular, $\varphi^\sharp_{\bar{0}}(f) = 0$, proving the claim. □

Proposition 4.22 (Products of finitely generated superspaces)**.** *Let X and Y be (locally) finitely generated. Then $X \times Y$ exists in the category of regular superspaces of countable girth, and is (locally) finitely generated.*

Proof. The question is local. So, passing to open subspaces, assume that embeddings $i : X \to \mathbb{A}^p \times \operatorname{Spec} A$ and $j : Y \to \mathbb{A}^r \times \operatorname{Spec} B$ are given, with the corresponding vanishing ideals $\mathcal{I}$ and $\mathcal{J}$. Clearly,

$$W \coloneqq \mathbb{A}^p \times \operatorname{Spec} A \times \mathbb{A}^r \times \operatorname{Spec} B \cong \mathbb{A}^{p+r} \times \operatorname{Spec}(A \otimes B)$$

exists in the category of regular superspaces. Let $U \subseteq \mathbb{A}^p_0 \times \mathbb{A}^r_0$ be open such that $(i_0 \times j_0)(X_0 \times Y_0)$ is closed in U, and $\mathcal{K}$ be the ideal of $\mathcal{O}_W|_U$ generated by $p_1^\sharp(\mathcal{I})$ and $p_2^\sharp(\mathcal{J})$. Define the $\mathbb{K}$-superspace $Z \coloneqq (Z_0, \mathcal{O}_Z)$ by $Z_0 \coloneqq X_0 \times Y_0$ and $\mathcal{O}_Z \coloneqq (i_0 \times j_0)^{-1}\big(\mathcal{O}_W|_U/\mathcal{K}\big)$, together with the canonical embedding $k : Z \to W$. Let $t : Z^\circ \to Z$ be the tidying of Z. Then $k^\circ \coloneqq k \circ t$ is an embedding and Z° is finitely generated by Corollary 4.20.

We have $(p_1 \circ k^\circ)^\sharp(\mathcal{I}) = 0$ and $(p_2 \circ k^\circ)^\sharp(\mathcal{J}) = 0$. Hence, the morphisms $p_1 \circ k^\circ$ and $p_2 \circ k^\circ$ factor uniquely through i and j, respectively, giving rise to morphisms $\pi_1 : Z^\circ \to X$ and $\pi_2 : Z^\circ \to Y$.

Let R be a regular superspace of countable girth, and $\phi : R \to X$, $\psi : R \to Y$ be morphisms. Since R is regular, $\varphi \coloneqq (i \circ \phi, j \circ \psi) : R \to W$ exists. Since $(p_1 \circ \varphi)^\sharp(\mathcal{I}) = 0$ and $(p_2 \circ \varphi)^\sharp(\mathcal{J}) = 0$, we have $\varphi^\sharp(\mathcal{K}) = 0$. Moreover, $\varphi_0(R_0) \subseteq k_0(Z_0)$, so φ factors uniquely through k, and since R has countable girth, it factors uniquely through k°, by Proposition 2.34. Denote the resulting morphism $R \to Z^\circ$ by ϱ.

We have $i \circ \pi_1 \circ \varrho = p_1 \circ k^\circ \circ \varrho = p_1 \circ \varphi = i \circ \phi$, so that $\pi_1 \circ \varrho = \phi$. Similarly, $\pi_2 \circ \varrho = \psi$. Let $\sigma : R \to Z^\circ$ be another morphism such that $\pi_1 \circ \sigma = \phi$ and $\pi_2 \circ \sigma = \psi$. Then $k^\circ \circ \sigma = (i \circ \phi, j \circ \psi) = k^\circ \circ \varrho$, and it follows that $\sigma = \varrho$. This proves the assertion. □

Corollary 4.23 (Limits of finitely generated superspaces)**.** *The category of locally finitely generated $\mathbb{K}$-superspaces is finitely complete.*

Proof. Since $\mathbb{A}^p$ is Hausdorff, any locally finitely generated $\mathbb{K}$-superspace is locally Hausdorff. Hence, the equaliser Z of morphisms $\phi, \psi : X \to Y$ locally finitely generated $\mathbb{K}$-superspaces exists in the category of $\mathbb{K}$-superspaces of countable girth, owing to Proposition 2.37.

By the same token, such a Z is tidy, and locally possesses embeddings into some $\mathbb{A}^p \times \operatorname{Spec} A$, since so does X. But these embeddings are necessarily tidy, by Corollary 2.33, so that Z is locally finitely generated. The assertion now follows from Proposition 4.22. □

The corollary allows to introduce the following notation.

Notation 4.24 (Fibre products and fibres)**.** Let $\varphi : X \to Z$ and $\psi : Y \to Z$ be morphisms of locally finitely generated superspaces. By virtue of Corollary 4.23, the fibred product $X \times_Z Y$ exists in the category of locally finitely generated superspaces.

In particular, if ψ is the inclusion of a point $z \in Z_0$ ($Y = *$ and $\psi_0(*) = z$), the fibre product is denoted by $\varphi^{-1}(z)$ or X_z. It is called the *fibre at z* of φ (or of X/Z).

We note another corollary to Proposition 4.22.

Corollary 4.25. *The reduction and body functors preserve finite limits in the category of locally finitely generated $\mathbb{K}$-superspaces.*

Proof. For the reduction functor, this is true by Proposition 4.21 and by the construction of equalisers and products (see the proofs of Proposition 2.37 and Proposition 4.22).

For the body functor, observe first that finite limits of even locally finitely generated $\mathbb{K}$-superspaces are again even, so the category of even locally finitely generated

$\mathbb{K}$-superspaces is finitely complete, in view of Corollary 4.23. Since the body functor is right adjoint to the inclusion of even locally finitely generated $\mathbb{K}$-superspaces into arbitrary locally finitely generated $\mathbb{K}$-superspaces, by Proposition 4.21 and Proposition 2.10, it preserves finite limits. □

We end this subsection by a discussion of morphisms and coordinates in the framework we have introduced. To treat the case where $\mathbb{K} \neq \Bbbk$ properly, we need a piece of notation. Consider the following super-vector space $V = V_{\bar{0}} \oplus V_{\bar{1}}$ over $(\mathbb{K}, \Bbbk)$: By definition, the odd part is $V_{\bar{1}} = \mathbb{K}^q$ and the even part is $V_{\bar{0}} = \mathbb{K}^p$, considered as a $\Bbbk$-vector space. We let $\mathbb{A}^{p|q}_{\mathbb{K}}$ be the affine superspace associated with V. If $\mathbb{K} = \Bbbk$, then $\mathbb{A}^{p|q}_{\mathbb{K}} = \mathbb{A}^{p|q}$, and if $\mathbb{K} \neq \Bbbk$, then $\mathbb{A}^{p|q}_{\mathbb{K}} = \mathbb{A}^{2p|q}$ as affine superspaces over $(\mathbb{K}, \Bbbk)$.

If $\mathbb{K} = \Bbbk$, then we denote the standard coordinate system on $\mathbb{A}^{p|q}_{\mathbb{K}} = \mathbb{A}^{p|q}$ by x_a, ξ_b. For the case of $\mathbb{K} \neq \Bbbk$, we let ξ_b, $b = 1, \dots, q$, be the standard coordinate functions coming from the $V_{\bar{1}}$ factor, and define x_a, y_a, $a = 1, \dots, p$, to be given by

$$\mathbb{C}^p \to \mathbb{R} : z = (x_1 + iy_1, \dots, x_p + iy_p) \mapsto \begin{cases} x_a, \\ y_a, \end{cases}$$

respectively. In what follows, it is understood that the y_a are only present for $\mathbb{K} \neq \Bbbk$.

Proposition 4.26. *Let X be a locally finitely generated $\mathbb{K}$-superspace. Then, for any non-negative integers p and q, there is a bijection*

$$\operatorname{Hom}(X, \mathbb{A}^{p|q}_{\mathbb{K}}) \to \Gamma\big(\mathcal{O}^p_{X,\bar{0}} \times \mathcal{O}^q_{X,\bar{1}}\big),$$

natural in X, given by

$$\varphi \mapsto \big(\varphi^\sharp(x_1 + iy_1), \dots, \varphi^\sharp(x_p + iy_p), \varphi^\sharp(\xi_1), \dots, \varphi^\sharp(\xi_q)\big).$$

The point of the proposition is that the tuples of superfunctions defining morphisms are not required to be $\Bbbk$-valued. In its *proof*, we use the following construction.

Lemma 4.27. *Let X be a $\mathbb{K}$-superspace and $\tilde{X} \coloneqq (X_0, \mathcal{O}_{X,\bar{0}})$. There is a canonical morphism $\psi_X : X \to \tilde{X}$, satisfying the following universal property: For any even $\mathbb{K}$-superspace Y and any morphism $\varphi : X \to Y$, there is a unique morphism $\tilde{\varphi} : \tilde{X} \to Y$ such that $\varphi = \tilde{\varphi} \circ \psi_X$. In particular, $X \mapsto \tilde{X}$ defines a functor and the ψ_X define a natural transformation $\mathrm{id} \to \tilde{\cdot}$. The functor $\tilde{\cdot}$ maps (open resp. closed) embeddings to (open resp. closed) embeddings and preserves the subcategory $\mathbf{SSp}^{\mathrm{lfg}}_{\mathbb{K}}$.*

Proof. Define $\psi = \psi_X : X \to \tilde{X}$ by setting $\psi_0 \coloneqq \mathrm{id}_{X_0}$ and $\psi^\sharp$ to be the natural sheaf map $\mathcal{O}_{X,\bar{0}} \to \mathcal{O}_X$. Let Y be an even $\mathbb{K}$-superspace. Any morphism $\varphi : X \to Y$ factors uniquely through ψ_X, because $\varphi^\sharp$ and all the superfunctions on Y are even. Thus, $X \mapsto \tilde{X}$ is the object map of a left adjoint to the inclusion of even $\mathbb{K}$-superspaces in $\mathbb{K}$-superspaces. The statements about embeddings are obvious, because of the graded nature of vanishing ideals.

Finally, let us show that $\tilde{X}$ is locally finitely generated if X is. Firstly, let A be the even part of the Grassmann algebra $\mathcal{O}_{\mathbb{A}^{0|M}}$. Then

$$A = \mathbb{K}\big[t_{ij} \;\big|\; 1 \leqslant i < j \leqslant M\big]/I\ , \quad \text{where} \quad I \coloneqq \big(t_{ij}t_{k\ell} \;\big|\; \{i,j\} \cap \{k,\ell\} \neq \varnothing\big).$$

This is an even Weil $\mathbb{K}$-algebra, and in case $X = \mathbb{A}^{N|M}$, we have $\tilde{X} = \mathbb{A}^N \times \operatorname{Spec} A$. Applying Corollary 4.18, we find that $\tilde{X}$ is finitely generated in case $X = \mathbb{A}^N$. Moreover, it follows from the definitions that $\tilde{X}$ is tidy if X is. Thus, according to the above statements about embeddings, Corollary 2.33 and the definitions, $\tilde{X}$ is indeed locally finitely generated if X is. □

Proof of Proposition 4.26. Unless $\mathbb{K} \neq \Bbbk$, there is nothing to prove, in view of Proposition 3.14. Thus, we may assume that $\mathbb{K} = \mathbb{C}$ and $\Bbbk = \mathbb{R}$. Moreover, since $\mathbb{A}^{p|q}_{\mathbb{K}} = \mathbb{A}^p_{\mathbb{K}} \times \mathbb{A}^{0|q}$, where $\mathbb{A}^p_{\mathbb{K}}$ is the affine $\mathbb{K}$-superspace associated with the $\mathbb{R}$-vector space $\mathbb{C}^p$, we may by the same token assume that $q = 0$.

Since $\mathbb{A}^p_{\mathbb{K}}$ is even, we may by Lemma 4.27 assume that X is even. In case $X = \mathbb{A}^N$, Corollary 4.3 shows that a morphism $X \to \mathbb{A}^p_{\mathbb{K}}$ is the same as a map $\mathbb{R}^N \to \mathbb{C}^p$ of class $\mathcal{C}^\varpi$. In turn, this is the same as a p-tuple of $\mathbb{C}$-valued functions on $\mathbb{R}^N$ of class $\mathcal{C}^\varpi$. This proves the assertion in case $X = \mathbb{A}^N$.

In general, we may assume that there is an embedding $j : X \to \mathbb{A}^N$. Since $\mathbb{A}^p_{\mathbb{K}} \cong \mathbb{A}^{2p}$, Proposition 4.10 shows that any morphism $X \to \mathbb{A}^p_{\mathbb{K}}$ locally factors through j. By the assertion for $X = \mathbb{A}^N$, and by Proposition 1.20, it follows that the map in the statement of the proposition is injective. Conversely, suppose we are given $\varphi_1, \dots, \varphi_p \in \Gamma(\mathcal{O}_X)$. Passing to an open cover, we may assume that $\varphi_a = j^\sharp(\psi_a)$ for some $\psi_a \in \Gamma(\mathcal{O}_{\mathbb{A}^N})$, $a = 1, \dots, N$. There is a morphism $\psi : \mathbb{A}^N \to \mathbb{A}^p_{\mathbb{K}}$ such that $\psi^\sharp(x_a + iy_a) = \psi_a$. Defining $\varphi : X \to \mathbb{A}^p_{\mathbb{K}}$ by $\varphi \coloneqq \psi \circ j$, we are done. □

We are now in a good position to discuss the meaning of morphisms from Weil thickened superspaces to a regular superspace with a global chart.

Scholium 4.28 (Morphisms and products)**.** Let S be a locally finitely generated $\mathbb{K}$-superspace, and A a Weil $\mathbb{K}$-superalgebra. Consider the Weil thickened superspace $S^A = S \times \operatorname{Spec} A$ with structure sheaf $\mathcal{O}_{S^A} = \mathcal{O}_S \otimes A$, defined in Construction 2.18. Then S^A is again locally finitely generated. Recall that $(S^A)_0 = S_0$. On one hand, we have

$$j^\sharp_{(S^A)_0}(f \otimes 1) = j^\sharp_{S_0}(f) \quad \text{for all} \quad f \in \mathcal{O}_S(U);$$

on the other hand, $j^\sharp_{(S^A)_0} = 0$ on $\mathcal{O}_S \otimes \mathfrak{m}$.

Let X be a regular $\mathbb{K}$-superspace with a global system of coordinates $x = (x_a)$. Then X is isomorphic to an open subspace of some $\mathbb{A}^{p|q}$. By Proposition 3.14, the set $X(S^A)$ of morphisms $\varphi : S^A \to X$ is in natural bijection with

$$\big\{ s = (s_a) \in (\Gamma(\mathcal{O}_S) \otimes A)^n \;\big|\; |s_a| = |x_a|\,,\; s(S_0) \subseteq x(X_0) \big\},$$

where $n \coloneqq p + q$. Under the bijection, φ maps to s if and only if $s = \varphi^\sharp(x)$.

Now, let $m \coloneqq k + \ell$, where $\dim \mathfrak{m} = k|\ell$, and choose a homogeneous basis $e^1, \dots, e^m$ of $\mathfrak{m}$. For any a, we may write

$$s_a = s^0_a \otimes 1 + \sum_{b=1}^m s^b_a \otimes e^b,$$

with unique $s^b_a \in \Gamma(\mathcal{O}_S)$, $b = 0, \dots, m$. Since $j^\sharp_{(S^A)_0}(s^b_a \otimes e^b) = 0$ for $b > 0$, we have

$$X(S^A) \cong \big\{ s = (s^b_a) \in \Gamma(\mathcal{O}_S)^{n\cdot(m+1)} \;\big|\; |s^b_a| = |x_a| + |e^b|, s^0(S_0) \subseteq x(X_0) \big\},$$

where we agree to write $e^0 \coloneqq 1$, and $s^b \coloneqq (s^b_a)_{a=1,\dots,n}$ for $b = 0, \dots, m$. Observe that here, the mapping condition only concerns s^0.

Applying Proposition 3.14, s^0 corresponds to a morphism $\varphi^0 : S \to X$, and by Proposition 4.26, s^b, for $b > 0$, corresponds to a morphism $\varphi^b : S \to \mathbb{A}^{p|q}_{\mathbb{K}}$ if $|e^b| = \bar{0}$, and to a morphism $\varphi^b : S \to \mathbb{A}^{q|p}_{\mathbb{K}}$ otherwise. The correspondences are given *via*

$$s^0 = \varphi^{0\sharp}(x) \quad \text{and} \quad s^b_a = \begin{cases} \varphi^{b\sharp}(z) & |e^b| = \bar{0}, \\ \varphi^{b\sharp}(\Pi z) & |e^b| = \bar{1}. \end{cases}$$

Here, $z = (x_a + iy_a, \xi_a)$ denotes the standard coordinate system of $\mathbb{A}^{p|q}_{\mathbb{K}}$, and Πz denotes the standard coordinate system of $\mathbb{A}^{q|p}_{\mathbb{K}}$.

In summary, we have for $\dim X = p|q$ and $\dim A = (k+1)|\ell$ a bijection

$$X(S^A) \cong X(S) \times \mathbb{A}^{p|q}_{\mathbb{K}}(S)^k \times \mathbb{A}^{q|p}_{\mathbb{K}}(S)^\ell = (X \times \mathbb{A}^{kp+\ell q|kq+\ell p}_{\mathbb{K}})(S) \tag{4.2}$$

which depends on the choice of a (global) coordinate system on X.

In this form, the bijection, however, also depends on the choice of a homogeneous basis of $\mathfrak{m}$; only the first component is entirely canonical.

To give a bijection, which only depends on the choice of the coordinate system (x_a) on X, consider the affine superspace $\mathfrak{m}^{p|q}$ associated with the $(\mathbb{K},\Bbbk)$-supervector space obtained by forgetting the $\mathbb{K}$-structure on the even part of $\mathfrak{m} \otimes_{\mathbb{K}} \mathbb{K}^{p|q}$. The data $(s^b_a)_{b>0}$ correspond to a morphism $\varphi^+ : S \to \mathfrak{m}^{p|q}$ *via* $s^b_a = \varphi^{+\sharp}(e_b \otimes z_a)$. We obtain a bijection

$$X(S^A) \cong X(S) \times \mathfrak{m}^{p|q}(S) = (X \times \mathfrak{m}^{p|q})(S), \tag{4.3}$$

where φ and (φ^0, φ^+) are related by

$$(\mathrm{id} \otimes \varepsilon)(\varphi^\sharp(f)) = \varphi^{0\sharp}(f) \quad \text{and} \quad (\mathrm{id} \otimes \mu)(\varphi^\sharp(z_a)) = \varphi^{+\sharp}(\mu \otimes z_a);$$

here, $\varepsilon : A \to \mathbb{K}$ is the unique algebra morphism, and $f \in \mathcal{O}_X(U)$, $\mu \in \mathfrak{m}^*$ are arbitrary. The obtained bijection is natural in S and the pairs $(X, (x_a))$.

4.3. **Relative supermanifolds.** In this section, we can finally introduce relative supermanifolds. In what follows, let S be a locally finitely generated $\mathbb{K}$-superspace. We denote by $\mathbf{SSp}^{\mathrm{lfg}}_S$ the full subcategory of $\mathbf{SSp}_S$ consisting of all X/S which are locally finitely generated as $\mathbb{K}$-superspaces. In the case of $S = *$, we write $\mathbf{SSp}^{\mathrm{lfg}}$.

In this, we are assuming that $\mathbb{K}$, $\Bbbk$ and ϖ are understood. If this is not the case, we will write $\mathbf{SSp}^{\varpi,\mathrm{lfg}}_{\mathbb{K},\Bbbk,S}$ and $\mathbf{SSp}^{\varpi,\mathrm{lfg}}_{\mathbb{K},\Bbbk}$, respectively.

Definition 4.29 (Relative affine superspace)**.** We let $\mathbb{A}^{p|q}_S \coloneqq S \times \mathbb{A}^{p|q}$, the product of locally finitely generated superspaces. Let $t = (t_a)$ be the standard coordinate system on $\mathbb{A}^{p|q}$. By abuse of notation, the collection $(p_2^\sharp(t_a))$ of superfunctions on $\mathbb{A}^{p|q}_S$ will again be denoted by $t = (t_a)$.

We call this system of superfunctions the *standard fibre coordinate system* and say it is in *standard order* if the original system of superfunctions on $\mathbb{A}^{p|q}$ was. As with coordinate systems, we take the liberty also to denote this by (t, θ) if we wish to see the parity explicitly.

Proposition 4.30 (Morphisms with relative affine target)**.** *Let X/S be a relative $\mathbb{K}$-superspace in $\mathbf{SSp}^{\mathrm{lfg}}_S$. There is a natural bijection*

$$\mathrm{Hom}_S\big(X, \mathbb{A}^{p|q}_S\big) \to \big\{(\varphi_a) \in \Gamma(\mathcal{O}_X)^{p+q} \;\big|\; \forall a : |\varphi_a| = |t_a|\big\} : \varphi \mapsto \varphi^\sharp(t).$$

Proof. Since X is regular, it suffices to remark that there is a natural bijection $\mathrm{Hom}_S(X, S \times Y) = \mathrm{Hom}(X, Y)$ for locally finitely generated $\mathbb{K}$-superspaces Y. □

Definition 4.31 (Fibre charts and fibre coordinate systems)**.** Let X/S be a relative $\mathbb{K}$-superspace in $\mathbf{SSp}^{\mathrm{lfg}}_S$ and $U \subseteq S$ an open subspace. By Proposition 1.18, $X \times_S U$ exists and is an open subspace of X. Let V be an open subspace of $X \times_S U$.

A *local fibre chart* (defined on V/U) is then by definition an open embedding $\varphi : V/U \to \mathbb{A}^{p|q}_U/U$. If $V = X$, then we call φ a *global fibre chart*. The tuple $p|q$ is called the *(graded) fibre dimension* of φ.

Let φ be a local fibre chart. The tuple $(x, \xi) \coloneqq \varphi^\sharp(t, \theta)$ (also denoted $x = (x_a) \coloneqq \varphi^\sharp(t_a)$) is called a system of *local fibre coordinates* (defined on V/U). If φ is a global chart, we call it a system of *global fibre coordinates*. Due to Proposition 4.30, fibre charts and fibre coordinate systems are in bijection. We will also call $p|q$ the *(graded) dimension* of (x, ξ).

Given a system of local fibre coordinates (x, ξ) defined on V/U, any $f \in \mathcal{O}_X(V_0)$ has a unique representation

$$f = \sum\nolimits_I f_I \xi^I, \quad \text{where} \quad \xi^I \coloneqq \xi_{i_1} \cdots \xi_{i_k} \tag{4.4}$$

and $f_I = x^\sharp(g_I)$ for some $g_I \in \mathcal{O}_{\mathbb{A}^p_U}(\varphi_0(V_0))$. Here, by abuse of notation, x denotes the morphism $V/U \to \mathbb{A}^p_U/U$ determined by $x^\sharp(t_j) = x_j$.

Proposition 4.30 and Definition 4.31 at once give the following.

Corollary 4.32 (Relative morphisms *vs.* fibre coordinates)**.** *Let X/S and Y/S be in $\mathbf{SSp}_S^{\mathrm{lfg}}$ and $y = (y_a)$ a global fibre coordinate system in standard order on Y. The following map is a bijection:*

$$\mathrm{Hom}_S(X, Y) \to \big\{ x \in \Gamma(\mathcal{O}_X)_{\bar{0}}^p \times \Gamma(\mathcal{O}_X)_{\bar{1}}^q \;\big|\; x(X_0) \subseteq y(Y_0) \big\} \colon \varphi \mapsto \varphi^\sharp(y).$$

Definition 4.33 (Relative supermanifolds)**.** Let X/S be in $\mathbf{SSp}_S^{\mathrm{lfg}}$. If, for every point $x \in X_0$, there exist an open subspace V/U of X/S such that $x \in V_0$ and a local fibre chart of X/S defined on V/U, then X/S is called a *super-premanifold* over S. If in addition, X_0 is a Hausdorff topological space, then X/S is called a *supermanifold over S*. In case $S = *$, X is simply called a *supermanifold*.

By reason of Corollary 4.3, the graded dimensions of all local fibre charts at $x \in X_0$ are identical. We denote their common value by $\dim_{S,x} X$ and call this the *(graded) fibre dimension at x*. For $S = *$, we simply write $\dim_x X$ and speak of the *(graded) dimension at x*. If $\dim_{S,x} X$ (resp. $\dim_x X$) is independent of x, then we say that X has *pure (fibre) dimension*.

The full subcategory of $\mathbf{SSp}_S$ whose objects are the supermanifolds over S is denoted by $\mathbf{SMan}_S$. For $S = *$, we write $\mathbf{SMan}$. If $\mathbb{K}$, $\Bbbk$, and ϖ are not understood, we write $\mathbf{SMan}^\varpi_{\mathbb{K},\Bbbk,S}$ and $\mathbf{SMan}^\varpi_{\mathbb{K},\Bbbk}$, respectively.

If S is a supermanifold, then in the literature [2, 7], supermanifolds over S have been previously considered, under the name of *families of supermanifolds*.

Lemma 4.34. *Let X/S be a super-premanifold over S. When S is a super-premanifold, then so is X, and $\dim_x X = \dim_s S + \dim_{S,x} X$, for $p_{X,0}(x) = s \in S_0$.*

Proof. The question is local, and $\mathbb{A}^{r|s} \times \mathbb{A}^{p|q} = \mathbb{A}^{p+r|q+s}$. □

Lemma 4.35. *Let X be a super-premanifold over S. Then X_0 resp. $X_{\bar{0}}$ is a super-premanifold over S_0 resp. $S_{\bar{0}}$. If $S = *$, then $X_0 = X_{\bar{0}}$.*

Proof. The statements are local, and for $\mathbb{A}^{p|q}_S$, they follow from Corollary 4.25. □

Proposition 4.36 (Gluing of supermanifolds)**.** *Let $(U_i, X_i/U_i, \varphi_{ij})$ be gluing data for super-premanifolds over S. The glued superspace X/S is a super-premanifold over S. Moreover, if S and the X_i are paracompact, then so is X.*

Proof. The existence of X/S follows from Proposition 1.20. Since we have isomorphisms $X_i/U_i \to (X \times_S U_i)/U_i$, it is clear that X possesses an open cover by local fibre charts, so that it is a super-premanifold over S.

To establish the paracompactness assertion, we drop the subscripts $(-)_0$ and work in the category of topological spaces. Let X be the space obtained by gluing the X_i, and $p : X \to S$ the projection. Let $\mathcal{V} = (V_j)$ be an open cover of X. Since p is open and S is paracompact, we may by passing to a refinement assume that (U_i) is locally finite and each $p(V_j)$ is contained in some U_i. This gives open covers $\mathcal{V}_i$ of X_i such that $\mathcal{V} = \bigcup_i \mathcal{V}_i$. Replacing each of the $\mathcal{V}_i$ by a locally finite open refinement, we obtain a locally finite open refinement of $\mathcal{V}$. Hence, X is paracompact. □

Proposition 4.37 (Base change)**.** *Let X/S be a super-premanifold over S and T/S be in $\mathbf{SSp}_S^{\mathrm{lfg}}$. Then $X \times_S T$ is a super-premanifold over T, called the* base change *of X/S. If X/S and T/S are supermanifolds over S, then $X \times_S T$ is a supermanifold over T and over S.*

Proof. For the first statement, it suffices to observe $\mathbb{A}_S^{p|q} \times_S T = \mathbb{A}_T^{p|q}$. As to the second, we remark that $X_0 \times_{S_0} T_0$ is Hausdorff in case X_0, T_0 are. □

Corollary 4.38. *Finite products in $\mathbf{SSp}_S^{\mathrm{lfg}}$ preserve $\mathbf{SMan}_S$.*

4.4. **Weil functors.** In this subsection, we construct the Weil functors, and relative versions thereof. These furnish a uniform framework for the study of (relative) even and odd tangent bundles, and their higher order versions. Originally, Weil functors were defined by A. Weil [11] in the case of ordinary manifolds. For supermanifolds, they were introduced by Balduzzi–Carmeli–Fioresi [1]. We show that they (and their relative versions) exist in the category $\mathbf{SSp}_{\mathbb{K}}^{\mathrm{lfg}}$ and arise as (relative) *inner homs* in this category. This point of view is very efficient, as was originally noticed by Kontsevich for the odd tangent bundle in his seminal paper [6].

Definition 4.39 (Relative inner homs)**.** Let $\mathbf{C}$ be a finitely complete category, S be an object. We write $\mathbf{C}_S$ for the category of objects X/S, Y/S over S (*i.e.* morphisms $p_X : X \to S$ and $p_Y : Y \to S$, respectively) and morphisms $\varphi : X/S \to Y/S$ over S (*i.e.* morphism $X \to Y$ in $\mathbf{C}$ which relate the given morphisms p_X and p_Y). The hom sets in this category will be denoted by $\mathrm{Hom}_S(X, Y)$.

Given X/S and Y/S, we define a functor $\underline{\mathrm{Hom}}_S(X, Y)$ on $\mathbf{C}_S$ with values in the category **Sets** of sets and maps by

$$\underline{\mathrm{Hom}}_S(X, Y)(T) \coloneqq \mathrm{Hom}_S(T \times_S X, Y)$$

for all T/S. An object Z of $\mathbf{C}_S$ representing $\underline{\mathrm{Hom}}_S(X, Y)$ is called a *relative inner hom* from X to Y and denoted by $\underline{\mathrm{Hom}}_S(X, Y)$. That is, there is a natural bijection

$$\mathrm{Hom}_S(T, \underline{\mathrm{Hom}}_S(X, Y)) = \mathrm{Hom}_S(T \times_S X, Y)$$

for all T/S. If it exists, $\underline{\mathrm{Hom}}_S(X, Y)$ is unique up to canonical isomorphism. In case $S = *$ is a terminal object, we just write $\underline{\mathrm{Hom}}(X, Y)$ and drop the 'relative'. Of course, for objects X, Y of $\mathbf{C}_S$, $\underline{\mathrm{Hom}}_S(X, Y)$ is just $\underline{\mathrm{Hom}}(X, Y)$ in $\mathbf{C}_S$.

Remark 4.40. This definition (for $S = *$) mimics the customary identification of maps of sets $f : T \times X \to Y$ and $g : T \to \mathrm{Maps}(X, Y)$, given by $f(t, x) = g(t)(x)$.

If S is a set and $p_X : X \to S$, $p_Y : Y \to S$ are sets over S, then $\underline{\mathrm{Hom}}_S(X, Y)$ is represented by the disjoint union

$$\coprod_{s \in S} \big(\{s\} \times \mathrm{Maps}(X_s, Y_s)\big),$$

where $X_s = p_X^{-1}(s)$ and $Y_s = p_Y^{-1}(s)$ are the fibres of X/S and Y/S, respectively.

Relative inner homs are functorial in several ways, as we now discuss.

Construction 4.41 (Functoriality of inner homs)**.** Assume that $\underline{\mathrm{Hom}}_S(X, Y)$ is representable in the category $\mathbf{C}_S$. If X'/S is such that $\underline{\mathrm{Hom}}_S(X', Y)$ exists, consider $f : X/S \to X'/S$. We define a morphism

$$\underline{\mathrm{Hom}}_S(f, Y) : \underline{\mathrm{Hom}}_S(X', Y) \to \underline{\mathrm{Hom}}_S(X, Y)$$

as follows: Let $\varphi \in \underline{\mathrm{Hom}}_S(X', Y)(T) = \mathrm{Hom}_S(T \times_S X', Y)$. Define

$$\underline{\mathrm{Hom}}_S(f, Y)(\varphi) \coloneqq \varphi \circ (\mathrm{id}_T \times_S f) \in \mathrm{Hom}_S(T \times_S X, Y) = \underline{\mathrm{Hom}}_S(X, Y)(T).$$

Since this construction is natural, it defines the required morphism $\underline{\mathrm{Hom}}_S(f, Y)$.

It is clear that $\underline{\mathrm{Hom}}_S(\mathrm{id}_X, Y) = \mathrm{id}$ and

$$\underline{\mathrm{Hom}}_S(f \circ f', Y) = \underline{\mathrm{Hom}}_S(f', Y) \circ \underline{\mathrm{Hom}}_S(f, Y),$$

for $f : X'/S \to X''/S$ such that $\underline{\mathrm{Hom}}_S(X'', Y)$ exists, so that $\underline{\mathrm{Hom}}_S(-, Y)$ is a contravariant functor on the full subcategory of $\mathbf{C}_S$, on which it is defined.

Similarly, for any Y'/S such that $\underline{\mathrm{Hom}}_S(X, Y')$ is representable, we define, for $g : Y/S \to Y'/S$, a morphism $\underline{\mathrm{Hom}}_S(X, g) : \underline{\mathrm{Hom}}_S(X, Y) \to \underline{\mathrm{Hom}}_S(X, Y')$ by setting, for any $\psi \in \underline{\mathrm{Hom}}_S(X, Y)(T)$

$$\underline{\mathrm{Hom}}_S(X, g)(\psi) \coloneqq g \circ \psi \in \mathrm{Hom}_S(T \times_S X, Y') = \underline{\mathrm{Hom}}_S(X, Y')(T).$$

One sees easily that $\underline{\mathrm{Hom}}_S(X, -)$ is a functor on its domain of existence in $\mathbf{C}_S$.

Finally, assume that $\underline{\mathrm{Hom}}_S(X, Y)$, $\underline{\mathrm{Hom}}_S(Y, Z)$ are representable in $\mathbf{C}_S$. We define a morphism

$$\circ = \circ_{X,Y,Z} : \underline{\mathrm{Hom}}_S(Y, Z) \times \underline{\mathrm{Hom}}_S(X, Y) \to \underline{\mathrm{Hom}}_S(X, Z),$$

called *inner circle*, as follows. For

$$\varphi \in \underline{\mathrm{Hom}}_S(X, Y)(T) = \mathrm{Hom}_T(T \times_S X, T \times_S Y),$$
$$\psi \in \underline{\mathrm{Hom}}_S(Y, Z)(T) = \mathrm{Hom}_T(T \times_S Y, T \times_S Z),$$

we obtain a composite $\psi \circ \varphi \in \underline{\mathrm{Hom}}_S(X, Z)(T)$ as

$$\psi \circ_{\mathbf{C}} \varphi \in \mathrm{Hom}_T(T \times_S X, T \times_S Z),$$

after identifying with $\underline{\mathrm{Hom}}_S(X, Z)(T)$. The inner circle $\circ$ is a natural transformation of (tri-)functors. It enjoys the obvious associative law.

Moreover, if $\underline{\mathrm{Hom}}_S(X, X)$ is representable, there is a morphism

$$1_X : S \to \underline{\mathrm{Hom}}_S(X, X)$$

over S, corresponding to the element $\mathrm{id}_X \in \mathrm{Hom}_S(X, X) = \underline{\mathrm{Hom}}_S(X, X)(S)$. (Recall that S is the tensor unit of $\times_S$.) It is a left and right identity for the inner circle. In particular, $\underline{\mathrm{Hom}}_S(X, X)$ is a monoid object in $\mathbf{C}_S$, whenever it exists.

Weil functors are fibre bundles in $\mathbf{SMan}_S$. The relevant definitions are as follows.

Definition 4.42 (Relative fibre bundles)**.** Let X/S and B/S be in $\mathbf{SSp}_S^{\mathrm{lfg}}$. Let I be the set of connected components of B_0, $F = (F_i)_{i \in I}$ a collection of locally finitely generated superspaces over S, and assume given a morphism $p : X \to B$, so that X lies over B. A *relative local trivialisation* of X/B (or of p) with typical fibre F consists of an open subspace $U \subseteq B$ and an isomorphism

$$\tau : U \times_S F \coloneqq \coprod_{i \in I} U_i \times_S F_i \to U \times_B X = p^{-1}(U)$$

over U, where $U_i \coloneqq U \cap i$. Denote by $\tau^F_{X/S}(U)$ the set of all relative local trivialisations (U, τ), where U is fixed. Then $\tau^F_{X/S}$ is a sheaf of sets on B_0.

A subsheaf $\mathcal{A} \subseteq \tau^F_{X/S}$ is called a *relative fibre bundle atlas* with typical fibre F if for every $b \in B_0$, there is an open subspace $U \subseteq B$ such that $x \in U_0$ and $\mathcal{A}(U) \neq \varnothing$. Two such atlases are called *equivalent* if their union is still an atlas.

We say that X/B is a *relative fibre bundle* (over S) with typical fibre F if there exists an atlas of local trivialisations $\mathcal{A}$ of X/B with typical fibre F. In case $S = *$, we simply say that X/B is a *fibre bundle*.

If X/B is a relative fibre bundle whose typical fibre $F = (F_i)_{i \in I}$ is a collection of supermanifolds over S, then X is a supermanifold over S, according to Corollary 4.38.

In the following, let $A = \mathbb{K} \oplus \mathfrak{m}$ be a Weil $\mathbb{K}$-superalgebra. Moreover, we fix $S \in \mathbf{SSp}_{\mathbb{K}}^{\mathrm{lfg}}$. We will write

$$\operatorname{Spec}_S A \coloneqq S^A = S \times \operatorname{Spec} A.$$

Proposition 4.43. *For any relative supermanifold X/S, the relative inner hom*

$$T_S^A X \coloneqq \underline{\operatorname{Hom}}_S(\operatorname{Spec}_S A, X)$$

is representable in the category $\mathbf{SSp}_S^{\mathrm{lfg}}$, and lies in the subcategory $\mathbf{SMan}_S$. Moreover, we have the following facts.

(i) *For $A = \mathbb{K}$, we have a natural isomorphism $T_S^{\mathbb{K}} X \to X$.*

(ii) *For a morphism $\varphi : A \to B$ of Weil $\mathbb{K}$-superalgebras, there is a canonical*

$$T_S^\varphi : T_S^A X \to T_S^B X,$$

and the assignment $\varphi \mapsto T_S^\varphi$ is functorial.

(iii) *For any $\varphi : A \to B$ as in* (ii)*, the T_S^φ are natural transformations of functors $T_S^A \to T_S^B$. That is, there is a commutative diagram*

$$\begin{array}{ccc} T_S^A X & \xrightarrow{T_S^A \psi} & T_S^A Y \\ {\scriptstyle T_S^\varphi}\downarrow & & \downarrow{\scriptstyle T_S^\varphi} \\ T_S^B X & \xrightarrow[T_S^B \psi]{} & T_S^B Y \end{array}$$

for every morphism $\psi : X/S \to Y/S$. Here, $T_S^A \psi = \underline{\operatorname{Hom}}_S(\operatorname{Spec}_S A, \psi)$.

(iv) *The unique morphisms*

$$\mathbb{K} \xrightarrow{\eta} A \xrightarrow{\varepsilon} \mathbb{K}$$

of $\mathbb{K}$-superalgebras induce via (ii) *canonical morphisms*

$$X \xrightarrow{s_X^A \coloneqq T_S^\eta} T_S^A X \xrightarrow{p_X^A \coloneqq T_S^\varepsilon} X$$

of $\mathbb{K}$-superspaces over S. Then s_X^A is a section of p_X^A, i.e. *$p_X^A \circ s_X^A = \mathrm{id}_X$.*

(v) *We may consider the superspace $T_S^A X$ as lying over X by the projection $p_X^A : T_S^A X \to X$ and under X by the canonical section $s_X^A : X \to T_S^A X$. Then the morphisms $T_S^A \psi : T_S^A X \to T_S^A Y$ are over and under ψ. Moreover, the projection and canonical section define natural transformations $p_X : T_S^{(-)} \to \mathrm{id}$ and $s_X : \mathrm{id} \to T_S^{(-)}$, so the morphisms T_S^φ from* (ii) *are over and under X.*

(vi) *With the morphism $p_X^A : T_S^A X \to X$ as the projection, $T_S^A X$ is a relative fibre bundle over X, whose typical fibre at $x \in X_0$ with $p|q = \dim_{S,x} X$ is $\mathfrak{m}^{p|q}$. Thus, if $\dim A = (r+1)|s$, $d = \dim_{\Bbbk} \mathbb{K}$, then*

$$\dim_{X,y} T_S^A X = d(pr + qs) | ps + qr \quad \textit{whenever} \quad \dim_{S,x} X = p|q \,,\, p_X^A(y) = x.$$

Before showing the representability of $T_S^A X$, we construct the natural morphisms.

Construction 4.44. Consider a morphism of Weil $\mathbb{K}$-superalgebras, $\varphi : A \to B$ (say). This induces a morphism

$$\operatorname{Spec} \varphi \coloneqq (*, \varphi) : \operatorname{Spec} B \to \operatorname{Spec} A,$$

which we promote to a morphism

$$\operatorname{Spec}_S \varphi : \operatorname{Spec}_S B \to \operatorname{Spec}_S A$$

over S. Now, assume that $\underline{\operatorname{Hom}}_S(\operatorname{Spec}_S A, X)$ and $\underline{\operatorname{Hom}}_S(\operatorname{Spec}_S B, X)$ are representable. Then we may define

$$T_S^\varphi \coloneqq \underline{\operatorname{Hom}}_S(\operatorname{Spec}_S \varphi, X) : T_S^A X \to T_S^B X$$

by simply applying Construction 4.41.

Proof of Proposition 4.43. We first assume that X is an open subspace of the relative affine superspace $\mathbb{A}_S^{p|q} = S \times \mathbb{A}^{p|q}$. Consider the tensor product $\mathfrak{m}^{p|q} = \mathfrak{m} \otimes_{\mathbb{K}} \mathbb{K}^{p|q}$.

Recall the facts discussed at length in Scholium 4.28. In particular, for $T \in \mathbf{SSp}_S^{\mathrm{lfg}}$, we have, by the obvious extension of Equation (4.3) to the relative case, a bijection

$$\mathrm{Hom}_S(T \times_S \mathrm{Spec}_S A, X) = \mathrm{Hom}_S(T \times \mathrm{Spec}\, A, X) \cong \mathrm{Hom}_S(T, X \times \mathfrak{m}^{p|q}),$$

which is natural in T/S. Thus, $\underline{\mathrm{Hom}}_S(\mathrm{Spec}_S A, X)$ exists, and is represented by the direct product $X \times \mathfrak{m}^{p|q}$.

The general case now follows by passing to an open cover, using the functoriality of $\underline{\mathrm{Hom}}_S(\mathrm{Spec}_S A, -)$ from Construction 4.41, and applying Proposition 1.20. By the same token, the assignment $X \mapsto T_S^A X$ defines a functor.

For $A = \mathbb{K}$, we note that $\mathrm{Spec}_S A = S$, so we have natural bijections

$$\underline{\mathrm{Hom}}_S(\mathrm{Spec}_S A, X)(T) = \mathrm{Hom}_S(T \times_S S, X) = \mathrm{Hom}_S(T, X)$$

for T/S, so that $X \cong T_S^{\mathbb{K}} X$ by the Yoneda Lemma. This proves item (i).

Item (ii) follows immediately by Construction 4.44. Similarly, (iii) is implied by the associativity of the inner circle, *v.* Construction 4.41. Item (iv) is a special case of (ii). For (v), the first part is a special case of (iii), and the second follows from

$$\varepsilon_B \circ \varphi = \varepsilon_A \quad \text{and} \quad \eta_B = \varphi \circ \eta_A$$

for any morphism $\varphi : A \to B$ of Weil $\mathbb{K}$-superalgebras. The second equation is a consequence of unitality, and the first is one of locality, which is automatic. (For the latter, one may argue as in the proof of Proposition 2.14.)

As for (vi), it is by the functoriality and (v) sufficient to consider an open subspace $X \subseteq \mathbb{A}_S^{p|q}$, which was treated in detail at the beginning of the proof. □

Definition 4.45 (Relative Weil functor)**.** The functor T_S^A which maps $X/S \mapsto T_S^A X$ and morphisms $\psi : X/S \to Y/S$ to morphisms $\psi^A \coloneqq T_S^A \psi : T_S^A X \to T_S^A Y$ over ψ is called the (relative) *A-Weil functor*. Endowed with the projection p_X^A, $T_S^A X$ is called the (relative) *A-Weil bundle* of X.

Example 4.46. Let $X \in \mathbf{SMan}_S$ and consider the algebra $A \coloneqq \mathbb{K}[\varepsilon]/(\varepsilon^2)$ of dual numbers. There is a natural bijection from $\mathrm{Hom}_S(T, T_S^A X)$ to the set of pairs

$$\left\{(\varphi, \delta) \in \mathrm{Hom}_S(T, X) \times \mathrm{Hom}_{p_{T,0}^{-1}\mathcal{O}_S}\left(\varphi_0^{-1}\mathcal{O}_X, \mathcal{O}_T\right) \;\middle|\; \delta(fg) = \varphi^\sharp(f)\delta(g) + \delta(f)\varphi^\sharp(g)\right\}$$

of morphisms $\varphi : T/S \to X/S$ and even derivations along φ, which are linear over $\mathcal{O}_S$. One easily verifies that the transition functions of $T_S^A X$ are given by first derivatives. Hence, $T_S^A X$ is a $\mathbb{K}$-vector bundle whose sheaf of sections is seen to be

$$\mathcal{T}_{X/S} \coloneqq \underline{\mathcal{D}er}_{p_{X,0}^{-1}\mathcal{O}_S}(\mathcal{O}_X, \mathcal{O}_X),$$

the *relative tangent sheaf* of X, where $\underline{\mathcal{D}er}$ is the sheaf of graded derivations. Moreover, if E is a $\mathbb{K}$-vector bundle over X, then by definition, its sheaf of sections is the unique $\mathcal{O}_X$-module sheaf $\mathcal{E}$ such that for every Y/X, we have

$$\mathrm{Hom}(Y, Y \times_X E) = \mathrm{Hom}(\mathcal{O}_Y, p_Y^*(\mathcal{E})).$$

Therefore, $T_S^A X$ is the relative tangent bundle of X.

Proposition 4.47. *The relative Weil functor T_S^A enjoys the following properties:*

(i) *T_S^A preserves finite fibre products over S, and there is a natural base change isomorphism*

$$R \times_S T_S^A X \cong T_R^A(R \times_S X).$$

(ii) *If U is an open subspace of X, then $T_S^A U$ is an open subspace of $T_S^A X$.*
(iii) *If B is a Weil $\mathbb{K}$-superalgebra, then there is a natural isomorphism*

$$T_S^{A\otimes B} = T_S^A \circ T_S^B.$$

In particular, we have a natural isomorphism $T_S^A \circ T_S^B \cong T_S^B \circ T_S^A$.

Proof. (i) Preservation of binary products follows from the natural bijection

$$\begin{aligned}\operatorname{Hom}_S\big(T, T_S^A(X \times_S Y)\big) &= \operatorname{Hom}_S\big(T \times_S \operatorname{Spec}_S A, X \times_S Y\big)\\ &= \operatorname{Hom}_S\big(T \times_S \operatorname{Spec}_S A, X\big) \times \operatorname{Hom}_S\big(T \times_S \operatorname{Spec}_S A, Y\big)\\ &= \operatorname{Hom}_S\big(T, T_S^A X\big) \times \operatorname{Hom}_S\big(T, T_S^A Y\big)\end{aligned}$$

after an application of Yoneda's Lemma. The case of the empty product is similar:

$$\operatorname{Hom}_S\big(T, T_S^A(S)\big) = \operatorname{Hom}_S\big(T \times_S \operatorname{Spec}_S A, S\big) = *,$$

so that $T_S^A(S) \cong S$. For the base change, we compute

$$\begin{aligned}\operatorname{Hom}_R\big(T, R \times_S T_S^A X\big) = \operatorname{Hom}_S\big(T, T_S^A X\big) &= \operatorname{Hom}_S\big(T \times_S \operatorname{Spec}_S A, X\big)\\ &= \operatorname{Hom}_R\big(T \times_R \operatorname{Spec}_R A, R \times_S X\big)\\ &= \operatorname{Hom}_R\big(T, T_R^A(R \times_S X)\big),\end{aligned}$$

where we recall $\operatorname{Spec}_S A = S \times \operatorname{Spec} A$, and the superspace T/R is considered as lying over S by the composition with R/S.

Item (ii) is clear by construction, and for (iii), we observe that

$$\operatorname{Spec} A \otimes B = \operatorname{Spec} A \times \operatorname{Spec} B$$

by Proposition 2.20, so

$$\begin{aligned}\operatorname{Hom}_S\big(T, T_S^{A\otimes B} X\big) &= \operatorname{Hom}\big(T \times_S \operatorname{Spec}_S A \times_S \operatorname{Spec}_S B, X\big)\\ &= \operatorname{Hom}\big(T \times_S \operatorname{Spec}_S A, T_S^B X\big) = \operatorname{Hom}_S(T, T_S^A(T_S^B X)),\end{aligned}$$

for any locally finitely generated $\mathbb{K}$-superspace T/S. □

The following lemma is a generalisation of Proposition 4.43 (vi). It shows the utility of the general point of view on Weil functors, but we will not be applying it.

Lemma 4.48. *Let $\varphi : A \to B$ be a morphism of Weil $\mathbb{K}$-superalgebras. If φ is surjective, then $T_S^\varphi X : T_S^A X \to T_S^B X$ is a relative fibre bundle with typical fibre $(\ker\varphi)^{p|q}$ at x, where $\dim_{S,x} X = p|q$. If φ is injective, then locally, T_S^φ is isomorphic to $(\mathrm{id}, 0) : T_S^A X \to T_S^A X \times (\mathfrak{m}_B/\varphi(\mathfrak{m}_A))^{p|q}$.*

Proof. Both statements are local. Let X be an open subspace of $\mathbb{A}_S^{p|q}$. Then $T_S^\varphi : X \times \mathfrak{m}_A^{p|q} \to X \times \mathfrak{m}_B^{p|q}$ is given by $T_S^A = \mathrm{id}_X \times \varphi^{p|q}$, where $\varphi^{p|q} = \varphi|_{\mathfrak{m}_A} \otimes \mathrm{id}_{\mathbb{K}^{p|q}}$. If φ is injective, then there is a linear isomorphism $\mathfrak{m}_B \cong \mathfrak{m}_A \times \mathfrak{m}_B/\varphi(\mathfrak{m}_A)$. If it is surjective, then is an isomorphism $\mathfrak{m}_A \to \ker\varphi \times \mathfrak{m}_B$. □

Remarkably, the natural transformations T_S^φ already exhaust the full set of natural transformations between (relative) Weil functors. In the setting of manifolds, this was already noticed by A. Weil [11].

Proposition 4.49. *Let A and B be Weil $\mathbb{K}$-superalgebras. Consider T_S^A and T_S^B as functors $\mathbf{SMan}_S \to \mathbf{SMan}_S$. The map $\varphi \mapsto T_S^\varphi$ is a bijection from the set of algebra morphisms $A \to B$ to the set of natural transformations $T_S^A \to T_S^B$, which are natural under base change in T/S.*

In the *proof*, we note the following lemma.

Lemma 4.50. *Any natural transformation $T_S^A \to T_S^B$ is uniquely determined by its value at $X = \mathbb{A}_S^1$.*

Proof. Indeed, let $\sigma, \tau : T^A_S \to T^B_S$ be natural. We first show that they are determined by their values on $\mathbb{A}^{1|1}_S$. So, assume that $\sigma_{\mathbb{A}^{1|1}_S} = \tau_{\mathbb{A}^{1|1}_S}$.

The naturality of σ and τ gives $\sigma_X = \tau_X$ for $X = \mathbb{A}^1_S$ and $X = \mathbb{A}^{0|1}_S$. Since T^A_S and T^B_S preserve fibre products over S, we obtain equality for $X = \mathbb{A}^{p|q}_S$, where p and q are arbitrary. By item (ii) of Proposition 4.47, this equality extends to the case of an open subspace X of $\mathbb{A}^{p|q}_S$, and thus, to an arbitrary supermanifold over S, in view of Proposition 1.20.

This proves $\sigma = \tau$ under the assumption of $\sigma_X = \tau_X$ for $X = \mathbb{A}^{1|1}_S$. Naturality and item (iii) of Proposition 4.47 show that $T^C_S \tau_X = \tau_{T^C_S X}$ for any Weil superalgebra C. In particular, for $C = \mathbb{K}[\tau]$ (τ being an odd indeterminate), we have $T^C_S \mathbb{A}^{1|0} = \mathbb{A}^{1|1}_S$, by Equation (4.3). Then $\tau_{\mathbb{A}^{1|1}_S} = T^C_S \tau_{\mathbb{A}^{1|0}_S}$, and this finally proves the assertion. □

Proof of Proposition 4.49. Let $A_\Bbbk \coloneqq \Bbbk \oplus \mathfrak{m}$, where we forget the $\mathbb{K}$-structure on the even part. Then $A_\Bbbk$ is a monoid in the symmetric monoidal category of finite-dimensional super-vector spaces over $(\mathbb{K}, \Bbbk)$.

Observe that $T^A(\mathbb{A}^1) = A_\Bbbk$ as supermanifolds over $(\mathbb{K}, \Bbbk)$. Thus, the structure morphisms $a, m : \mathbb{A}^1 \times \mathbb{A}^1 \to \mathbb{A}^1$ of addition and multiplication, respectively, of the $\Bbbk$-algebra $\Bbbk$, give rise to operations $T^A(a)$ and $T^A(m)$ on $A_\Bbbk$. By construction of T^A, these are the same as the given addition and multiplication of $A_\Bbbk$. In particular, they are $\mathbb{K}$-linear resp. $\mathbb{K}$-bilinear on $\mathfrak{m}$.

By the base change formula in Proposition 4.47, we have $T^A_S(\mathbb{A}^1_S) = S \times T^A(\mathbb{A}^1)$, $T^A_S(a) = \mathrm{id}_S \times T^A(a)$, and $T^A_S(m) = \mathrm{id}_S \times T^A(m)$. Any natural transformation $T^A_S \to T^B_S$ gives rise to a morphism $\psi : S \times A \to S \times B$ over S, which commutes with the promotion of a and m to operations on $S \times A$ and $S \times B$. Naturality, applied to the embedding $\eta^A_{\mathbb{A}^1} : \mathbb{A}^1_S \to S \times A$, gives the unitality of ψ.

Finally, by the assumption of naturality, ψ is of the form $\mathrm{id}_S \times \varphi$ for some morphism $\varphi : A_\Bbbk \to B_\Bbbk$ of monoids in the category of super-vector spaces over $(\mathbb{K}, \Bbbk)$. Obviously, this extends uniquely to an even $\mathbb{K}$-superalgebra morphism $A \to B$. By Lemma 4.50, this proves our claim. □

References

[1] L. Balduzzi, C. Carmeli, and R. Fioresi, *The local functors of points of supermanifolds*, Expo. Math. **28** (2010), no. 3, 201–217, DOI 10.1016/j.exmath.2009.09.005.

[2] P. Deligne and J. W. Morgan, *Notes on supersymmetry*, Quantum Fields and Strings: A Course for Mathematicians, 1999, pp. 41–98.

[3] M. Demazure and P. Gabriel, *Groupes algébriques. Tome I: Géométrie algébrique, généralités, groupes commutatifs*, Masson & Cie, Éditeur, Paris, 1970.

[4] A. Grothendieck and J. A. Dieudonné, *Éléments de géométrie algébrique. I.*, Grundlehren der mathematischen Wissenschaften, vol. 166, Springer-Verlag, Berlin, 1971.

[5] I. Kolář, P. W. Michor, and J. Slovák, *Natural Operations in Differential Geometry*, Springer-Verlag, Berlin, 1993.

[6] M. Kontsevich, *Deformation quantization of Poisson manifolds*, Lett. Math. Phys. **66** (2003), no. 3, 157–216, DOI 10.1023/B:MATH.0000027508.00421.bf.

[7] D. A. Leites, *Introduction to the theory of supermanifolds*, Uspekhi Mat. Nauk **1** (1980), 3–57 (Russian); English transl., Russian Math. Surveys **35** (1980), no. 1, 1–64.

[8] Y. I. Manin, *Gauge Field Theory and Complex Geometry*, 2nd ed., Grundlehren der Mathematischen Wissenschaften, vol. 289, Springer-Verlag, Berlin, 1997.

[9] I. Moerdijk and G. E. Reyes, *Models for Smooth Infinitesimal Analysis*, Springer-Verlag, New York, 1991.

[10] H. Hohnhold, M. Kreck, S. Stolz, and P. Teichner, *Differential forms and 0-dimensional supersymmetric field theories*, Quantum Topol. **2** (2011), no. 1, 1–41, DOI 10.4171/QT/12.

[11] A. Weil, *Théorie des points proches sur les variétés différentiables*, Géométrie différentielle. Colloques Internationaux du Centre National de la Recherche Scientifique, Strasbourg, 1953, Centre National de la Recherche Scientifique, Paris, 1953, pp. 111–117 (French).

Universität zu Köln, Mathematisches Institut, Weyertal 86-90, 50931 Köln, Germany
E-mail address: alldridg@math.uni-koeln.de

Universität Paderborn, Institut für Mathematik, 33095 Paderborn, Germany
E-mail address: hilgert@math.upb.de

Institut É. Cartan, Université de Lorraine et C.N.R.S., 57045 Metz, France
Current address: Fakultät für Mathematik, Ruhr-Universität Bochum, 44780 Bochum, Germany
E-mail address: tilmann.wurzbacher@univ-lorraine.fr